

ISSN 2518-170X (Online)

ISSN 2224-5278 (Print)



«ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ
АКАДЕМИЯСЫ» РҚБ

«ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫ» РҚБ

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

РОО «НАЦИОНАЛЬНОЙ
АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН»

N E W S

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF
KAZAKHSTAN

SERIES

OF GEOLOGY AND TECHNICAL SCIENCES

4 (465)

JULY – AUGUST 2024

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of geology and technical sciences scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of geology and technical sciences in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of geology and engineering sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы «ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы» ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді геология және техникалық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАНПК сообщает, что научный журнал «Известия НАНПК. Серия геологии и технических наук» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАНПК. Серия геологии и технических наук в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по геологии и техническим наукам для нашего сообщества.

Бас редактор

ЖҰРЫНОВ Мұрат Жұрынұлы, химия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, «Қазақстан Республикасы Ұлттық ғылым академиясы» РҚБ-нің президенті, АҚ «Д.В. Сокольский атындағы отын, катализ және электрохимия институтының» бас директоры (Алматы, Қазақстан) **Н = 4**

Ғылыми хатшы

АБСАДЫКОВ Бахыт Нарикбайұлы, техника ғылымдарының докторы, профессор, ҚР ҰҒА жауапты хатшысы, А.Б. Бектұров атындағы химия ғылымдары институты (Алматы, Қазақстан) **Н = 5**

Редакциялық алқа:

ӘБСАМЕТОВ Мәліс Құдысұлы (бас редактордың орынбасары), геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, «У.М. Ахмедсафина атындағы гидрогеология және геоэкология институтының» директоры (Алматы, Қазақстан) **Н = 2**

ЖОЛТАЕВ Герой Жолтайұлы (бас редактордың орынбасары), геология-минералогия ғылымдарының докторы, профессор, Қ.И. Сатпаев атындағы геология ғылымдары институтының директоры (Алматы, Қазақстан) **Н = 2**

СНОУ Дэниел, Ph.D, қауымдастырылған профессор, Небраска университетінің Су ғылымдары зертханасының директоры (Небраска штаты, АҚШ) **Н = 32**

ЗЕЛЬТМАН Реймар, Ph.D, табиғи тарих мұражайының Жер туралы ғылымдар бөлімінде петрология және пайдалы қазбалар кен орындары саласындағы зерттеулердің жетекшісі (Лондон, Англия) **Н = 37**

ПАНФИЛОВ Михаил Борисович, техника ғылымдарының докторы, Нанси университетінің профессоры (Нанси, Франция) **Н = 15**

ШЕН Пин, Ph.D, Қытай геологиялық қоғамының тау геологиясы комитеті директорының орынбасары, Американдық экономикалық геологтар қауымдастығының мүшесі (Пекин, Қытай) **Н = 25**

ФИШЕР Аксель, Ph.D, Дрезден техникалық университетінің қауымдастырылған профессоры (Дрезден, Берлин) **Н = 6**

КОНТОРОВИЧ Алексей Эмильевич, геология-минералогия ғылымдарының докторы, профессор, РҒА академигі, А.А. Трофимука атындағы мұнай-газ геологиясы және геофизика институты (Новосибирск, Ресей) **Н = 19**

АГАБЕКОВ Владимир Енокович, химия ғылымдарының докторы, Беларусь ҰҒА академигі, Жаңа материалдар химиясы институтының құрметті директоры (Минск, Беларусь) **Н = 13**

КАТАЛИН Стефан, Ph.D, Дрезден техникалық университетінің қауымдастырылған профессоры (Дрезден, Берлин) **Н = 20**

СЕЙТМҰРАТОВА Элеонора Юсуповна, геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА корреспондент-мүшесі, Қ.И. Сатпаев атындағы Геология ғылымдары институты зертханасының меңгерушісі (Алматы, Қазақстан) **Н = 11**

САҒЫНТАЕВ Жанай, Ph.D, қауымдастырылған профессор, Назарбаев университеті (Нұр-Сұлтан, Қазақстан) **Н = 11**

ФРАТТИНИ Паоло, Ph.D, Бикокк Милан университеті қауымдастырылған профессоры (Милан, Италия) **Н = 28**

«ҚР ҰҒА» РҚБ Хабарлары. Геология және техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктеуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан Республикасының Ақпарат және қоғамдық даму министрлігінің Ақпарат комитетінде 29.07.2020 ж. берілген № **KZ39VPU00025420** мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *геология, мұнай және газды өңдеудің химиялық технологиялары, мұнай химиясы, металдарды алу және олардың қосындыларының технологиясы.*

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекен-жайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., тел.: 272-13-19

<http://www.geolog-technical.kz/index.php/en/>

© «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ, 2024

Главный редактор

ЖУРИНОВ Мурат Журинович, доктор химических наук, профессор, академик НАН РК, президент РОО «Национальной академии наук Республики Казахстан», генеральный директор АО «Институт топлива, катализа и электрохимии им. Д.В. Сокольского» (Алматы, Казахстан) **Н = 4**

Ученый секретарь

АБСАДЫКОВ Бахыт Нарикбаевич, доктор технических наук, профессор, ответственный секретарь НАН РК, Институт химических наук им. А.Б. Бектурова (Алматы, Казахстан) **Н = 5**

Редакционная коллегия:

АБСАМЕТОВ Малис Кудысович, (заместитель главного редактора), доктор геологоминералогических наук, профессор, академик НАН РК, директор Института гидрогеологии и геоэкологии им. У.М. Ахмедсафина (Алматы, Казахстан) **Н = 2**

ЖОЛТАЕВ Герой Жолтаевич, (заместитель главного редактора), доктор геологоминералогических наук, профессор, директор Института геологических наук им. К.И. Сатпаева (Алматы, Казахстан) **Н=2**

СНОУ Дэниел, Ph.D, ассоциированный профессор, директор Лаборатории водных наук университета Небраски (штат Небраска, США) **Н = 32**

ЗЕЛЬТМАН Реймар, Ph.D, руководитель исследований в области петрологии и месторождений полезных ископаемых в Отделе наук о Земле Музея естественной истории (Лондон, Англия) **Н = 37**

ПАНФИЛОВ Михаил Борисович, доктор технических наук, профессор Университета Нанси (Нанси, Франция) **Н=15**

ШЕН Пин, Ph.D, заместитель директора Комитета по горной геологии Китайского геологического общества, член Американской ассоциации экономических геологов (Пекин, Китай) **Н = 25**

ФИШЕР Аксель, ассоциированный профессор, Ph.D, технический университет Дрезден (Дрезден, Берлин) **Н = 6**

КОНТОРОВИЧ Алексей Эмильевич, доктор геолого-минералогических наук, профессор, академик РАН, Институт нефтегазовой геологии и геофизики им. А.А. Трофимука СО РАН (Новосибирск, Россия) **Н = 19**

АГАБЕКОВ Владимир Енокович, доктор химических наук, академик НАН Беларуси, почетный директор Института химии новых материалов (Минск, Беларусь) **Н = 13**

КАТАЛИН Стефан, Ph.D, ассоциированный профессор, Технический университет (Дрезден, Берлин) **Н = 20**

СЕЙТМУРАТОВА Элеонора Юсуповна, доктор геолого-минералогических наук, профессор, член-корреспондент НАН РК, заведующая лабораторией Института геологических наук им. К.И. Сатпаева (Алматы, Казахстан) **Н=11**

САГИНТАЕВ Жанай, Ph.D, ассоциированный профессор, Назарбаев университет (Нурсултан, Казахстан) **Н = 11**

ФРАТТИНИ Паоло, Ph.D, ассоциированный профессор, Миланский университет Бикокк (Милан, Италия) **Н = 28**

«Известия РОО «НАН РК». Серия геологии и технических наук».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и общественного развития Республики Казахстан № **KZ39VPY00025420**, выданное 29.07.2020 г.

Тематическая направленность: *геология, химические технологии переработки нефти и газа, нефтехимия, технологии извлечения металлов и их соединений.*

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, оф. 219, тел.: 272-13-19

<http://www.geolog-technical.kz/index.php/en/>

© РОО «Национальная академия наук Республики Казахстан», 2024

Editorial chief

ZHURINOV Murat Zhurinovich, doctor of chemistry, professor, academician of NAS RK, president of the National Academy of Sciences of the Republic of Kazakhstan, general director of JSC “Institute of fuel, catalysis and electrochemistry named after D.V. Sokolsky» (Almaty, Kazakhstan) **H = 4**

Scientific secretary

ABSADYKOV Bakhyt Narikbaevich, doctor of technical sciences, professor, executive secretary of NAS RK, Bekturov Institute of chemical sciences (Almaty, Kazakhstan) **H = 5**

Editorial board:

ABSAMETOV Malis Kudysovich, (deputy editor-in-chief), doctor of geological and mineralogical sciences, professor, academician of NAS RK, director of the Akhmedsafin Institute of hydrogeology and hydrophysics (Almaty, Kazakhstan) **H=2**

ZHOLTAEV Geroy Zholtaevich, (deputy editor-in-chief), doctor of geological and mineralogical sciences, professor, director of the institute of geological sciences named after K.I. Satpayev (Almaty, Kazakhstan) **H=2**

SNOW Daniel, Ph.D, associate professor, director of the laboratory of water sciences, Nebraska University (Nebraska, USA) **H = 32**

ZELTMAN Reyman, Ph.D, head of research department in petrology and mineral deposits in the Earth sciences section of the museum of natural history (London, England) **H = 37**

PANFILOV Mikhail Borisovich, doctor of technical sciences, professor at the Nancy University (Nancy, France) **H=15**

SHEN Ping, Ph.D, deputy director of the Committee for Mining geology of the China geological Society, Fellow of the American association of economic geologists (Beijing, China) **H = 25**

FISCHER Axel, Ph.D, associate professor, Dresden University of technology (Dresden, Germany) **H=6**

KONTOROVICH Aleksey Emilievich, doctor of geological and mineralogical sciences, professor, academician of RAS, Trofimuk Institute of petroleum geology and geophysics SB RAS (Novosibirsk, Russia) **H = 19**

AGABEKOV Vladimir Enokovich, doctor of chemistry, academician of NAS of Belarus, honorary director of the Institute of chemistry of new materials (Minsk, Belarus) **H = 13**

KATALIN Stephan, Ph.D, associate professor, Technical university (Dresden, Berlin) **H = 20**

SEITMURATOVA Eleonora Yusupovna, doctor of geological and mineralogical sciences, professor, corresponding member of NAS RK, head of the laboratory of the Institute of geological sciences named after K.I. Satpayev (Almaty, Kazakhstan) **H=11**

SAGINTAYEV Zhanay, Ph.D, associate professor, Nazarbayev University (Nursultan, Kazakhstan) **H = 11**

FRATTINI Paolo, Ph.D, associate professor, university of Milano-Bicocca (Milan, Italy) **H = 28**

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: RPA «National Academy of Sciences of the Republic of Kazakhstan» (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Social Development of the Republic of Kazakhstan **No. KZ39VPY00025420**, issued 29.07.2020.

Thematic scope: *geology, chemical technologies for oil and gas processing, petrochemistry, technologies for extracting metals and their connections.*

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, Almaty, 050010, tel. 272-13-19

<http://www.geolog-technical.kz/index.php/en/>

© National Academy of Sciences of the Republic of Kazakhstan, 2024

NEWS of the National Academy of Sciences of the Republic of Kazakhstan

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224–5278

Volume 4. Number 466 (2024), 117–130

<https://doi.org/10.32014/2024.2518-170X.429>

UDC 528.88, 626.81

© G.M. Iskaliyeva^{1,2*}, N.K. Sydyk¹, A.K. Kalybayeva^{1,3}, M.S. Sagat¹, A. Samat⁴, 2024

¹ Institute of Ionosphere, Almaty, Kazakhstan;

²Kazakh National Agrarian University, Almaty, Kazakhstan;

³Al-Farabi Kazakh National University, Almaty, Kazakhstan;

⁴Xinjiang Institute of Ecology and Geography, Urumqi, China.

E-mail: igm.ionos@gmail.com

USE OF WATER INDICES FOR WATERBODIES IN THE ESIL WATER MANAGEMENT BASIN

Iskaliyeva Gulnara — Leading researcher at the Laboratory of Space and Geoinformation Technologies, Institute of Ionosphere, Ph.D. student, Kazakh National Agrarian Research University, Almaty, Kazakhstan

E-mail: igm.ionos@gmail.com; <https://orcid.org/0000-0002-3183-728X>;

Sydyk Nurmakhambet — Head of laboratory at the Laboratory of Space and Geoinformation Technologies, Institute of Ionosphere, Ph.D. student, Al Farabi Kazakh National University, Almaty, Kazakhstan

E-mail: nurmahambet.s@gmail.com; <https://orcid.org/0000-0003-1429-2393>;

Kalybayeva Aigerim — Senior Researcher at the Laboratory of Space Monitoring of Emergency Situations, Ph.D. student, Al Farabi Kazakh National University, Almaty, Kazakhstan

E-mail: kalybayeva.aigerim@gmail.com; <https://orcid.org/0000-0001-6088-6876>;

Sagat Madina — Engineer at the Laboratory of Space and Geoinformation Technologies, Institute of Ionosphere, Almaty, Kazakhstan

E-mail: sagat.madina1@gmail.com; <https://orcid.org/0009-0004-0174-5677>;

Alim Samat — PhD, researcher in Research Centre for Ecology and Environment of Central Asia, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi, 830011, China

E-mail: alim_smt@ms.xjb.ac.cn; <https://orcid.org/0000-0002-9091-6033>.

Abstract. According to the historical information, there are more than 48 thousand large and small lakes in Kazakhstan, a significant part of which are located in the north of the republic. However, limnological studies of most lakes in Kazakhstan are still quite weak. This study examined the applicability of Landsat (8,9) and Sentinel-2 satellite images for the detection of surface water bodies in the Esil water management basin (WMB) using water indices: NDVI, AWEI, AWEIsh, WRI, NDWI, MNDWI, NDMI, NDTI, NDSI. The MNDWI and AWEI indices were found to be the best for water distribution in the two test lakes of this study. NDTI showed the worst results. There are 831 lakes in the Esil River basin, of which only 455 lakes have a surface area of more than 1 km² and 71 lakes have a surface area of more than 10 km². According to an analysis based on various indices, in 2023, 770 lakes were discovered in the Esil River basin, of which only 452 lakes have a surface area of more than 1 km² and 70 lakes have a surface area of more than 10 km². Thus, there was a decrease in the number of 149 lakes with a surface area of more than 1 km² and 27 lakes with a surface area of more than 10 km² by compared to the data file.

Keywords: Water indices, Remote sensing, Esil water management basin, GIS; lakes, water surfaces

Acknowledgements. This research was funded by the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. BR18574092). This research was implemented with the focus on the partial requirements of PhD Candidate – Iskaliyeva Gulnara’s doctoral dissertation on the «Assessment of the dynamics of the lake systems of the Esil water basin based on remote sensing data using GIS technology» at Kazakh National Agrarian Research University, Almaty, Kazakhstan.

© Г.М. Искалиева^{1,2*}, Н.Қ. Сыдық¹, А.К. Қалыбаева^{1,3}, М.С. Сағат¹,
А. Самат⁴, 2024

¹ Ионосфера институты, Алматы, Қазақстан;

²Қазақ ұлттық аграрлық университеті, Алматы, Қазақстан;

³Әл-Фараби атындағы Қазақ ұлттық университеті, Алматы, Қазақстан;

⁴Шыңжаң экология және география институты, Үрімші, Қытай.

E-mail: igm.ionos@gmail.com

ЕСІЛ СУ ШАРУАШЫЛЫҒЫ БАССЕЙНІНДЕГІ СУ ОБЪЕКТИЛЕРІ ҮШІН СУ ИНДЕКСТЕРІН ПАЙДАЛАНУ

Искалиева Гульнара — ғарыштық және геоақпараттық технологиялар зертханасының жетекші ғылыми қызметкері, Ионосфера институты, Қазақ ұлттық аграрлық зерттеу университетінің докторанты, Алматы, Қазақстан

E-mail: igm.ionos@gmail.com; <https://orcid.org/0000-0002-3183-728X>;

Сыдық Нұрмахамбет — ғарыштық және геоақпараттық технологиялар зертханасының меңгерушісі, Ионосфера институты, әл-Фараби атындағы Қазақ ұлттық университетінің докторанты, Алматы, Қазақстан

E-mail: nurmahambet.s@gmail.com; <https://orcid.org/0000-0003-1429-2393>;

Қалыбаева Әйгерім — төтенше жағдайлардың ғарыштық мониторингі зертханасының аға ғылыми қызметкері, Ионосфера институты, әл-Фараби атындағы Қазақ ұлттық университетінің докторанты, Алматы қ., Қазақстан

E-mail: kalybayeva.aigerim@gmail.com; <https://orcid.org/0000-0001-6088-6876>;

Сағат Мадина — ғарыштық және геоақпараттық технологиялар зертханасының инженері, Ионосфера институты, Алматы, Қазақстан

E-mail: sagat.madinal@gmail.com; <https://orcid.org/0009-0004-0174-5677>;

Әлім Самат — философия докторы, Орталық Азияның экология және қоршаған орта ғылыми-зерттеу орталығының ғылыми қызметкері, Шыңжаң экология және география институты, Қытай ғылым академиясының, Үрімші, 830011, Қытай

E-mail: alim_smt@ms.xjb.ac.cn; <https://orcid.org/0000-0002-9091-6033>.

Аннотация. Мұрағаттық деректер бойынша Қазақстан Республикасы аймағында 48 мыңнан астам үлкен және кіші көлдер орналасқан және олардың айтарлықтай бөлігі республиканың трансшекаралық солтүстік бөлігінде орналасқан. Дегенмен, еліміздің көптеген көлдерінің лимнологиялық зерттелуі әлі де әлсіз. Бұл еңбектегі зерттеу NDVI, AWEI, AWEIsh, WRI, NDWI, MNDWI, NDMI, NDTI, NDSI су индекстерін Landsat (8,9) және Sentinel-2 ғарыштық суреттеріне қолданыла отырып, Есіл су шаруашылығы бассейніндегі жер үсті су объектілерін анықтау үшін жүргізілді. Осы зерттеу бойынша қарастырылған көлдерде су бетін анықтау үшін қолданылған индекстерге қарағанда MNDWI және AWEI индекстері жақсы нәтиже беретіні анықталды, ал ең қолайсыз индекс болып төмен нәтижелі NDTI болды. Бұрынғы

зерттеу мәліметтерде көсетілгендей, 2013 жылы Есіл су шаруашылығы бассейнінде 831 көл болған, оның тек 455 көлінің ауданы 1 км²-ден асады және 71 көл – ауданы 10 км²-ден асқан. Әр түрлі индекстер комбинациясы негізінде талдау бойынша 2023 жылы зерттеу аумағында 770 көл идентификацияланды, оның ішінде: ауданы 1 км²-ден асатын 452 көл және ауданы 10 км²-ден асатын 70 көл анықталды. Осылайша, мұрағаттық деректермен салыстырғанда, жалпы ауданы 1 км²-ден асатын көлдер 149 көлге, ал ауданы 10 км²-ден асатын объектілер саны бойынша 27 көлге азайды.

Түйін сөздер: су индекстері, ЖҚЗ, Есіл су шаруашылық бассейні, ГАЖ, көлдер, су беті

© Г.М. Искалиева^{1,2*}, Н.К. Сыдык¹, А.К. Калыбаева^{1,3}, М.С. Сагат¹, А. Самат⁴,
2024

¹ Институт ионосферы, Алматы, Казахстан;

² Казахский национальный аграрный университет, Алматы, Казахстан;

³ Казахский национальный университет им. аль-Фараби, Алматы, Казахстан;

⁴ Сельскохозяйственный Университет, Пловдив, Болгария.

E-mail: igm.ionos@gmail.com

ИСПОЛЬЗОВАНИЕ ВОДНЫХ ИНДЕКСОВ ДЛЯ ВОДНЫХ ОБЪЕКТОВ В ЕСИЛЬСКОМ ВОДОХОЗЯЙСТВЕННОМ БАССЕЙНЕ

Искалиева Гульнара — ведущий научный сотрудник лаборатории космических и геоинформационных технологий, Институт ионосферы, докторант Казахского национального аграрного исследовательского университета, Алматы, Казахстан

E-mail: igm.ionos@gmail.com; <https://orcid.org/0000-0002-3183-728X>;

Сыдык Нурмахамбет — заведующий лабораторией космических и геоинформационных технологий Института ионосферы, докторант Казахского национального университета имени аль-Фараби, Алматы, Казахстан

E-mail: nurmahambet.s@gmail.com; <https://orcid.org/0000-0003-1429-2393>;

Калыбаева Айгерим — старший научный сотрудник лаборатории космического мониторинга чрезвычайных ситуаций, Институт ионосферы, докторант Казахского национального университета имени аль-Фараби, Алматы, Казахстан

E-mail: kalybayeva.aigerim@gmail.com; <https://orcid.org/0000-0001-6088-6876>;

Сагат Мадина — инженер лаборатории космических и геоинформационных технологий, Институт ионосферы, Алматы, Казахстан

E-mail: sagat.madinal@gmail.com; <https://orcid.org/0009-0004-0174-5677>;

Алим Самат — доктор философии, научный сотрудник Научно-исследовательского центра экологии и окружающей среды Центральной Азии, Синьцзянский институт экологии и географии, Китайская академия наук, Урумчи, 830011, Китай

E-mail: alim_smt@ms.xjb.ac.cn; <https://orcid.org/0000-0002-9091-6033>.

Аннотация. По архивным данным, на территории Казахстан насчитывается более 48 тысяч больших и малых озер, их основная часть находится в северной части республики. При этом большинство озер Казахстана по лимнологическим показателям слабо изучено. В этом исследовании изучалась применимость водных индексов: NDVI, AWEI, AWEIsh, WRI, NDWI, MNDWI, NDMI, NDTI, NDSI для снимков со спутников Landsat (8,9) и Sentinel-2 по обнаружению поверхностных водных объектов в Есильском водохозяйственном бассейне (ВХБ). Было обнаружено,

что для выделения воды на тестовых озерах данного исследования индексы MNDWI и AWEI подходят лучше, чем остальные. В то же время индекс NDTI показал наихудшие результаты. По архивным данным, в 2013 г. в Есильском ВХБ было 831 озеро, из которых 455 озер имело площадь более 1 км² и 71 озеро были с площадью больше 10 км². На основе комбинации различных индексов на территории Есильского ВХБ за 2023 год обнаружено 770 озеро. По анализу было обнаружено с площадью более 1 км² 452 озер и 70 озеро – свыше 10 км². Таким образом, по сравнению с архивными данными, произошло сокращение на 149 озеро с площадью свыше 1 км² и на 27 озер с площадью более 10 км².

Ключевые слова: водные индексы, ДЗЗ, Есильский водохозяйственный бассейн, ГИС, озера, водная поверхность

Introduction

In modern conditions, considering the growing global shortage of natural water, the issue of rational use, protection, and restoration of lake resources becomes especially relevant. Inefficient use and improper management of water reserves often lead to the depletion of water resources and the reduction of the inland lakes area.

Desertification and salinization in arid areas significantly contribute to the reduction of lake areas. The size of water bodies affects the climate of adjacent regions, influencing air humidity levels and, consequently, the nature of precipitation in the lake basin.

In Kazakhstan, according to archival data, there are more than 48,000 large and small lakes, a significant part of which are located in the territory of interest. Kazakhstan possesses a total of 48,262 lakes, with 45,248 being categorized as small lakes covering an area less than 1 km², 296 classified as large lakes spanning over 10 km², and 21 exceeding 100 km² in size. The latter group represents 60 % of the total water surface area of lakes in Kazakhstan (Frumin & Krashanovskaya, 2010).

Lake water is used in various industries, but its use is currently not well-organized (Imanaliyev et al., 2022). Therefore, a comprehensive assessment and inventory of the lake fund's current state in Kazakhstan is necessary, using modern methodologies and combining all available data in a cartographic format.

The limnological study of most lakes in Kazakhstan, their condition, and water quality and quantity is still quite weak. Studies of the lakes of Kazakhstan, initiated in the 1950s by the Geography Sector of the Academy of Sciences of the Kazakh SSR, were discontinued in the 1990s. Thus, there has been a break of more than 25 years in lake monitoring. During this time, due to changes in natural character and anthropogenic factors, many lakes have changed their hydrological, morphometric, and hydrochemical characteristics (Yang et al., 2015).

Such alterations in the presence of surface water structures have a profound impact on both agricultural and industrial productivity (Sharma et al., 2015; Anderson et al., 2012), in addition to affecting environmental and ecological security (Palmer et al., 2015; Yamazaki et al., 2015; Du et al., 2016; Liu et al., 2015).

Data on the lake locations, their volumes, and water quality, as well as conditions and possibilities for their rational use, are outdated and cannot objectively address water security issues in the Republic.

Remote sensing is an expeditiously advancing technology that can offer cheap and dependable information regarding environmental modifications at local, regional, and global levels due to the prolonged accumulation of repetitive data (Lee et al., 2018). For instance,

Pekel et al. (2016) utilized all Landsat images to identify alterations in surface water reservoirs over the previous three decades on a worldwide scale. Several techniques for water extraction have been devised and implemented for images acquired through remote sensing.

McFeeters formulated the most rudimentary spectral index for water detection, known as the NDWI (McFeeters et al., 1996), by utilizing the green and NIR bands of the Landsat (TM) image to optimize the identification of water. Furthermore, McFeeters proposed that setting a threshold of 0 enables the differentiation between water and the surrounding background using NDWI (9).

Xu substituted the NIR band within McFeeters' NDWI with the SWIR band and developed a modified NDWI (MNDWI) (Xu et al., 2006). Water distribution in urban regions was delineated based on the MNDWI index utilizing Landsat (ETM+) imagery, revealing that MNDWI outperforms NDWI in discerning water from shadows.

Feyisa et al. (2014) introduced the Automatic Water Extraction Index (AWEI) for the identification of water bodies, featuring two components: AWEI_{sh} is tailored to eliminate shadow pixels, while AWEI_{ns} is targeted towards regions with urban characteristics. Furthermore, a multitude of other indices have been developed in various research endeavors for diverse sensors and locations, each exhibiting varying levels of accuracy (Malahlela et al., 2016; Fisher et al., 2013).

Study area. According to our country's Water Code, the basin principle of water resource management is in effect. There are 8 water management basins: Zhaiyk-Caspian, Aral-Syr Darya, Tobol-Turgay, Esil, Nura-Sarysu, Chu-Talas, Balkhash-Alakol, and Irtysh. The water resources within each basin are hydrologically connected, so it is expedient to consider the water management situation in the country in terms of river basins. The Yesil Water Basin, located in the northern and central regions of Kazakhstan, covers a vast area spanning the Akmola, Kostanay, and North Kazakhstan regions (Figure 1). Fed primarily by the Yesil River, the basin receives its water from numerous significant tributaries originating in the Kokshetau Hills in the north and the Ulytau Ranges in the south. The region's largely flat topography, combined with the presence of various closed basins, creates favorable conditions for surface water retention and the development of lake reservoirs. The basin covers an area of approximately 176,000 square kilometers and has experienced declining water runoff for many years, during which much of the meltwater is used to fill basins and floodplains in the river channels. As a result, runoff volumes decrease, primarily due to snow accumulation in the river bed, resulting in a reduction in the area of the active catchment basin. Such hydrological patterns are usually characterized by dry conditions similar to those of a desert. Conversely, in years of high water levels, the relief exceeds its capacity, which facilitates the diversion of water into the main river bed and leads to a significant expansion of the active area of the catchment basin. The rapid onset of spring facilitates the unimpeded flow of meltwater into river channels, minimizing water losses and leading to high floods even with small snow reserves.

This determines the specific hydrological conditions of this region, characterized by extreme unevenness of runoff both in the long-term perspective and throughout the year.

In the Esil Water Management Basin, there are numerous lakes that are extremely important in the ecosystem of the region.

The lakes of the Esil Water Management Basin face a range of environmental problems. Climate change and unsustainable water use contribute to increased salinity and decreased water levels in lakes, threatening biodiversity and the sustainability of ecosystems.

The adverse impact on lake water quality caused by agricultural practices and industrial discharges leads to pollution and further deterioration of plant and animal habitats. Extensive irrigation methods, leading to decreased water levels in lakes, have a detrimental effect on the ecological state of these water bodies.

The climate of the Esil Water Management Basin is continental, characterized by sharp seasonal and daily temperature fluctuations, low precipitation, and significant evaporation. Typical January temperatures range from -15 °C to -20 °C. The winter season is characterized by low temperatures and long snowfalls. The average July temperature ranges from +18 °C to +22 °C. The summer months are characterized by high temperatures and dry weather, with occasional precipitation. Annual precipitation ranges from 300 to 400 mm. Snow cover is established in November and lasts until March-April. Snowmelt is an important source of water replenishment in the spring. Significant daily and seasonal temperature fluctuations are a characteristic feature of the region's climate.




Figure 1 – Esil Water Management Basin (Esil WMB) Territory

The average air humidity throughout the year varies, reaching its lowest values in the summer and highest in the winter. High evaporation rates, especially in the summer months, contribute to the drying up of water bodies and decreased groundwater levels.

The climate conditions of the Esil Water Management Basin are the main factors of the formation and distribution of lakes in the region. Spring flooding, caused by snowmelt, is the main source of replenishment for rivers and lakes. Summer droughts and high evaporation lead to decreased water levels in water bodies and increased salinity.

Materials and methodology

The National Aeronautics and Space Administration (NASA, Washington, D.C., USA) launched the first Landsat satellite in 1972, and since then we have successfully launched Landsat 8–9 satellites are equipped with Operational Land Imager (OLI) or Operational Land Imager-2 (OLI-2) sensors, which increase the spectral resolution of the TM group by adding deep blue and cirrus bands. We analyzed and downloaded Landsat images to obtain water data from the website USGS Earthexplorer.

Sentinel-2 MSI images were downloaded from the Copernicus browser website, which serves as a repository for Earth observation satellite imagery from the European Space Agency. The image consists of 13 Sentinel-2A spectral bands, each with a different spatial resolution. Specifically, there are three visible bands and one near-infrared band with 10 m spatial resolution, a quad band dedicated to vegetation analysis, two short-wave near-infrared bands with 20 m resolution, and three thermal infrared bands with 60 m resolution. However, this study only focused on the use of blue (10 m), green (10 m), near-infrared (10 m), and short-wave infrared (20 m) due to their effectiveness in trapping water. data. We performed spatial sampling to compensate for the 10 m spatial resolution, taking into account the difference in spatial resolution between the SWIR and green/NIR bands. For the analysis, satellite images were used from September-October, a period without snow and when cloudiness did not exceed 10 %.

Data preprocessing stages included cropping of the study area, radiometric correction, and atmospheric correction. Landsat image values are stored as digital numbers representing electromagnetic radiation detected by the satellite's corresponding sensors. The purpose of radiometric correction is to convert the brightness value of a gray image into a bright glow of something comparable to the values of images that have undergone the same level of temporal or sensory correction. Atmospheric correction assumes that the overall brightness of an object on the ground measured by the sensor is not the true surface brightness, which includes error in the amount of radiation caused by atmospheric absorption, especially dispersion. Atmospheric correction is the process of eliminating errors caused by radiation from weather effects and ensuring the true surface brightness of objects on the ground.

Methodology of calculation of auxiliary indices.

To develop an algorithm for water surface extraction from LANDSAT (8,9) and Sentinel-2 satellite data, different spectral indices selected from the analyses of world experience were tested:

The first step was to calculate the Normalized Difference Vegetation Index to determine the boundaries of the lake:

$$\begin{aligned} \text{NDVI} &= (\text{NIR} - \text{RED})/(\text{NIR} + \text{RED}) \\ \text{Landsat} &= (\text{B5}-\text{B4})/(\text{B5}+\text{B4}) \\ \text{Sentinel 2} &= (\text{B8}-\text{B4})/(\text{B8}+\text{B4}) \end{aligned} \quad (1)$$

NDVI is defined by values from -1.0 to 1.0; here negative values occur mainly for clouds, water and snow.

AutomatedWaterExtractionIndex is calculated using the following formula, and water has a for this index positive value:

$$\begin{aligned}
 \text{AWEI} &= 4 * (\text{Green} - \text{SWIR2}) - (0.25 * \text{NIR} + 2.75 * \text{SWIR1}) \\
 \text{Landsat} &= 4*(\text{B3}-\text{B7})-(0.25*\text{B5}+2.75*\text{B6}) \\
 \text{Sentinel 2} &= 4*(\text{B3}-\text{B12})-(0.25*\text{B8}+2.75*\text{B11})
 \end{aligned} \tag{2}$$

The AWEI (Automated Water Extraction Index) index is specifically designed to mask non-aquatic pixels in urban environments. It is most effective on images with less than 15 % cloud cover and a modified AWEIsh index, designed to more accurately identify water in shaded areas and other dark surfaces. However, errors may occur on the classification screen in areas with highly reflective surfaces such as ice and snow.

$$\begin{aligned}
 \text{AWEIsh} &= \text{Blue} + 2.5 * \text{Green} - 1.5 * (\text{NIR} + \text{SWIR1}) - 0.25 * \text{SWIR2} \\
 \text{Landsat} &= \text{B2} + 2.5 * \text{B3} - 1.5 * (\text{B5} + \text{B6}) - 0.25 * \text{B7} \\
 \text{Sentinel 2} &= \text{B2} + 2.5 * \text{B3} - 1.5 * (\text{B8} + \text{B11}) - 0.25 * \text{B12}
 \end{aligned} \tag{3}$$

Water Ratio Index (WRI) is used to determine the moisture content of vegetation and is calculated as follows:

$$\begin{aligned}
 \text{WRI} &= (\text{GREEN} + \text{RED}) / (\text{NIR} + \text{SWIR2}) \\
 \text{Landsat} &= (\text{B3} + \text{B4}) / (\text{B5} + \text{B7}) \\
 \text{Sentinel 2} &= (\text{B3} + \text{B4}) / (\text{B8} + \text{B12})
 \end{aligned} \tag{4}$$

In the process of determining the indices, we obtain raster data about the water surface, where the pixel values will be 0 or higher; and values from 1 indicate bodies of water or objects containing moisture. This approach makes it possible to identify water systems in the study area.

The Normalized Difference Moisture Index (NDWI) is a relative index used to estimate the moisture content of vegetation, particularly in relation to the absorption of solar radiation. Thanks to its algorithm, this indicator facilitates a qualitative assessment of the level of humidity in the crown of plants. In contrast to the normalized vegetation index Differential (NDVI), NDWI index is sensitive to fluctuations in humidity and is also less sensitive to weather conditions. It is used to determine surface water in wetlands and to measure the availability of surface water. The typical range for lush vegetation is 0.1 to 0.4. This indicator is especially useful when the mid-infrared channel is not available.

The NDWI index is calculated as follows:

$$\begin{aligned}
 \text{NDWI} &= (\text{NIR} - \text{SWIR2}) / (\text{NIR} + \text{SWIR2}) \\
 \text{Landsat} &= (\text{B5} - \text{B7}) / (\text{B5} + \text{B7}) \\
 \text{Sentinel 2} &= (\text{B8} - \text{B12}) / (\text{B8} + \text{B12})
 \end{aligned} \tag{5}$$

Modified Normalized Difference Water Index (MNDWI) is used in various situations to improve the identification and monitoring of water bodies using remote sensing data. The index provides effective attenuation and eventual elimination of spectral interference from surfaces, soil and vegetation. It outperforms NDWI results, especially in scenarios where the studied coastline is very close to a man-made structure. As a result, determining the boundaries of water bodies in such cases provides higher accuracy without taking into account negative factors that interfere with work. MNDWI is particularly useful for accu-

rately identifying water bodies in the context of landscape and plant systems. His formula:

$$\begin{aligned} \text{MNDWI} &= (\text{Green} - \text{SWIR2}) / (\text{Green} + \text{SWIR2}) \\ \text{Landsat} &= (\text{B3} - \text{B7}) / (\text{B3} + \text{B7}) \\ \text{Sentinel 2} &= (\text{B3} - \text{B12}) / (\text{B3} + \text{B12}) \end{aligned} \quad (6)$$

The Normalized Differential Moisture Index (NDMI) is designed to monitor drought conditions and estimate fuel moisture content in wildfire-prone areas. The use of near-infrared (NIR) and short-wave infrared (SWIR) spectral bands provides connectivity while reducing the effects of interference and weather conditions, and improves the display of humidity levels using near-infrared (NIR) and short-wave bands. Spectral bands of infrared waves (SWIR), which are assigned a positive numerical value to the water content and measured using a special mathematical expression:

$$\begin{aligned} \text{NDMI} &= (\text{NIR} - \text{SWIR 1}) / (\text{NIR} + \text{SWIR 1}) \\ \text{Landsat} &= (\text{B5} - \text{B6}) / (\text{B5} + \text{B6}) \\ \text{Sentinel 2} &= (\text{B8} - \text{B11}) / (\text{B8} + \text{B11}) \end{aligned} \quad (7)$$

Standard algorithm (NDTI): index based on water turbidity. This indicator characterizes a decrease in water transparency due to impurities of inorganic and organic substances in suspension, as well as the development of plankton in the lake.

NDTI is calculated:

$$\begin{aligned} \text{NDTI} &= (\text{RED} - \text{GREEN}) / (\text{RED} + \text{GREEN}) \\ \text{Landsat} &= (\text{B4} - \text{B3}) / (\text{B4} + \text{B3}) \\ \text{Sentinel 2} &= (\text{B4} - \text{B3}) / (\text{B4} + \text{B3}) \end{aligned} \quad (8)$$

This index is suitable for identifying objects with actual humidity. To calculate the NDVI and NDTI indices, the characteristic quality of vegetation is determined; This demonstrates a pattern: the higher the NDVI, the greater the NDTI. The indices are useful for drought forecasting.

The Standardized Difference Snow Index (NDSI) was developed to detect the neutral layer of snow in clouds in Landsat TM images (bands 2 and 5). The index serves to reduce the impact of atmospheric conditions. In the case of snow, the NDSI value is usually greater than 0.4 and its calculation is based on:

$$\begin{aligned} \text{NDSI} &= (\text{GREEN} - \text{SWIR1}) / (\text{GREEN} + \text{SWIR1}) \\ \text{Landsat} &= (\text{B3} - \text{B6}) / (\text{B3} + \text{B6}) \\ \text{Sentinel 2} &= (\text{B3} - \text{B11}) / (\text{B3} + \text{B11}) \end{aligned} \quad (9)$$

The brightness of the snow layer makes it difficult to distinguish snow from clouds in images. Snow not only reflects light well in the visible part of the electromagnetic spectrum, but also absorbs it well in the near-infrared and short-wave parts of the spectrum. When exposed to solar radiation with a wavelength of 1.6 microns, snow becomes darker in color than clouds.

Results and discussion


The study evaluates the performance of water-dominated indices: NDVI, AWEI, AWEISH, WRI, NDWI, MNDWI, NDMI, NDTI, Landsat (8.9) NDSI satellite imagery and Sentinel-2 for the Yesil Basin . The analysis includes an examination of standards and ideal thresholds for defining surface waters specific to grasslands and forests. In conclusion, we implemented and evaluated a new approach that integrates indices and cross-sections of a study area as a function of elevation to improve the detection of surface water across the entire area. Figures 2 and 3 show an example of the results of calculating various water surface indices.




Sentinel-2 image for October 19, 2023




PlanetScope image for October 27, 2023.




NDVI Index




AWEI Index




AWEIsh Index




WRI Index




NDWI Index




MNDWI Index



NDMI Index



NDTI Index



NDSI Index

Figure 2 – Calculation of different water surface definition indices at Lake Kishi Karaoi


The study found that MNDWI and AWEI were better at identifying the water surface, while NDTI performed poorest results.




Sentinel-2 image for October 19, 2023




PlanetScope image for October 27, 2023.




NDVI Index



AWEI Index



AWEIsh Index



WRI Index




Figure 3 – Calculation of different indices of water surface definition at Lake Qalibek

Each of these indices has its own advantages and disadvantages. Using the NDVI index, you can clearly identify an area with cloudless, clear water that contrasts well with the background of clouds, ice and snow. This index typically identifies water, very wet soil, recently burned areas, and cloud shadows. The water surface with shallow water and vegetation is little differentiated. NDSI is mostly fully developed in winter or early spring, but is also useful for eliminating water surface class cloud shadows during snow-free periods.

Using archival data, the Esil Basin was found to have 9,812 lakes (Filonets et al., 1974), with 601 lakes larger than 1 km² and 97 lakes larger than 10 km². Satellite data processing in 2013 identified 831 lakes in the Esil Basin, with 455 lakes larger than 1 km² and 71 lakes larger than 10 km². Analysis in 2023 identified 770 lakes, with 452 lakes larger than 1 km² and 70 lakes larger than 10 km². This represents a reduction of 149 lakes larger than 1 km² and 27 lakes larger than 10 km² compared to archival data.

As a result of a study in which all the indices discussed above were used, according to the file, there are 9812 lakes in the Esil WMB (Filonets et al., 1974), of which 601 have an area of 1 km² and the surface area of 97 lakes is more than 10 km². According to the processing of satellite data, in 2013, 831 lakes were digitized in the Esil WMB, of which only 455 have a mirror area of more than 1 km² and 71 of which have a mirror area of more than 10 km². According to an analysis based on various indicators, in 2023, 770 lakes were

discovered on the territory of the Esil naval base, of which only 452 have a surface area of more than 1 km², and the surface of 70 lakes. Thus, according to the data of the array, there are 149 lakes with a surface area of more than 1 km², and there has been a decrease in 27 lakes with a surface area of more than 10 km².

The number of small lakes (up to 1 km²) is variable for the lowland regions of central, western and northern Kazakhstan, depending on the annual humidity in a particular region. The number of lakes in river deltas also varies.

The methodology was based on methods used to solve similar problems around the world. Numerous authors (Pekel et al., 2016) have used satellite and multispectral data to detect and monitor changes in water bodies, demonstrating the successful use of remote sensing in water resource management. The study was carried out using satellite altimetry, but the accuracy of the measurements depends on the size of the reservoir, surrounding topography, surface waves and winter ice cover. This method has performed poorly in most lakes larger than 10 km² (Busker, 2019; Smith et al., 2015).

Conclusion

This study examined the applicability of Landsat (8,9) and Sentinel-2 image indices to detect surface water bodies in steppe and forested areas using different water levels and compared the results.

By comparing the results of different water release rates under different conditions, we found that MNDWI and AWEI performed better than others in terms of water release, while NDTI performed worse in the two lakes presented in this study. The results also show that Landsat imagery (8,9) in combination with data from other Landsat sensors and other medium-resolution satellites generally allows for the study of changes in water regimes over long periods of time.

According to the results, it was revealed that the number of lakes from 1 to 10 km² varies quite a bit from year to year. Given the aridity of the study area, the response to climate change was observed mainly in lakes of small area. Warming will lead not only to an increase in air temperature, but also in the temperature of surface waters, which will increase the flow of moisture from the surface of the lake into the atmosphere. Analysis of the results confirms the high degree of variability of surface waters; This does not allow us to accurately define a group of lakes classified by surface.

This data set of lake volumes helps improve our current understanding of lake behavior, their representation in hydrologic models, and hence river basin modeling. It will also improve projections of river basin development in terms of climate change under different management scenarios and improve estimates of evaporation from open water bodies.

The Esil water management basin plays an important role in providing water and protecting the region's ecosystems. The lakes of the basin are important natural objects that require constant care and protection. Joint efforts by the state, the scientific community and the public can help protect and restore these unique water bodies, ensuring sustainable development and environmental stability.

REFERENCES

- Anderson M.C., Allen R.G., Morse A., Kustas W.P. (2012). Use of Landsat thermal imagery in monitoring evapotranspiration and managing water resources, *Remote Sens of Environment*. — 122:50–65. DOI: 10.1016/j.rse.2011.08.025 (in Eng.)
- Busker T., de Roo A., Gelati E., Schwatke C., Adamovic M., Bisselink B., Pekel J.-F., Cottam A. (2019). A global lake and lake volume analysis using a surface water dataset and satellite altimetry, *Hydrology and Earth System Sciences*. — 23:669–690. DOI:10.5194/hess-23-669-2019 (in Eng.)
- Du Y., Zhang Y., Ling F., Wang Q., Li W., Li X. (2016 a). Water Bodies' Mapping from Sentinel-2 Imagery with Modified Normalized Difference Water Index at 10-m Spatial Resolution Produced by Sharpening the SWIR Band, *Remote Sensing*. — 8:354. DOI: 10.3390/rs8040354 (in Eng.)
- Feyisa G.L., Meilby H., Fensholt R., Proud S.R. (2014). Automated water extraction index: A new technique for surface water mapping using Landsat imagery, *Remote Sensing of Environment*, — 140:23–35. DOI: 10.1016/j.rse.2013.08.029 (in Eng.)
- Filonets P.P., Omarov T.R. (1974). Lakes of Northern, Western and Eastern Kazakhstan. *Gidrometeoizdat, Russia*.
- Fisher A., Danaher T.A. (2013). A water index for SPOT 5 HRG satellite imagery, New South Wales, Australia, determined by linear discriminant analysis, *Remote Sensing*, — 5:5907. DOI: 10.3390/rs5115907 (in Eng.)
- Frumin G.T., Krashanovskaya Yu.V. (2014). Trophic status of lakes in Kazakhstan [Trophicheskiy status ozer Kazakhstana], *Terra Humana*, — 3:32 (in Russ.)
- Imanaliyev T., Koybakov S., Karlykhanov O., Amanbayeva B. & Bakiyev M. (2022). Prospects for the development of water resources management in the south of Kazakhstan, *News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences*. — 6(456):80–95. DOI:10.32014/2518-170X.240 (in Eng.)
- Lee J.K., Acharya T.D., Lee D.H. (2018). Exploring land cover classification accuracy of Landsat 8 image using spectral index layer stacking in hilly region of South Korea, *Sensors and Materials*, — 30:1–15. DOI: 10.18494/SAM.2018.1934 (in Eng.)
- Liu J., Zhang Y., Yuan D., Song X. (2015). Empirical Estimation of Total Nitrogen and Total Phosphorus Concentration of Urban Water Bodies in China Using High Resolution IKONOS Multispectral Imagery, *Water*. — 7:6551–6573. DOI: 10.3390/w7116551 (in Eng.)
- Malahlela O.E. (2016). Inland waterbody mapping: Towards improving discrimination and extraction of inland surface water features, *International Journal of Remote Sensing*. — 37:4574–4589. DOI: 10.1080/01431161.2016.1217441 (in Eng.)
- McFeeters S.K. (1996). The use of the Normalized Difference Water Index (NDWI) in the delineation of open water features, *International Journal of Remote Sensing*. — 17:1425–1432. DOI: 10.1080/01431169608948714 (in Eng.)
- Palmer S.C.J., Kutser T., Hunter P.D. (2015). Remote sensing of inland waters: Challenges, progress and future directions, *Remote Sensing of Environment*, — 157:1–8. DOI: 10.1016/j.rse.2014.09.021 (in Eng.)
- Pekel J.F., Cottam A., Gorelick N., Belward A.S. (2016). High-resolution mapping of global surface water and its long-term changes, *Nature*. — 540:418–422. DOI: 10.1038/nature20584 (in Eng.)
- Sharma R., Tateishi R., Hara K., Nguyen L. (2015). Developing Superfine Water Index (SWI) for Global Water Cover Mapping Using MODIS Data, *Remote Sensing*, — 7:13807–13841. DOI: 10.3390/rs71013807 (in Eng.)
- Smith L.C., Pavelsky T.M. (2015). Monitoring global surface water changes using multi-source satellite data, *Remote Sensing of Environment*. — 157:112–123. DOI:10.1016/j.rse.2014.06.008 (in Eng.)
- Xu H. (2006) Modification of normalised difference water index (NDWI) to enhance open water features in remotely sensed imagery, *International Journal of Remote Sensing*, 27:3025-3033. DOI: 10.1080/01431160600589179 (in Eng.)
- Yamazaki D., Trigg M.A., Ikeshima, D. (2015) Development of a global ~90m water body map using multi-temporal Landsat images, *Remote Sensing of Environment*, 171:337-351. DOI: 10.1016/j.rse.2015.10.014 (in Eng.)
- Yang Y., Liu Y., Zhou M., Zhang S., Zhan W., Sun C., Duan Y. (2015). Landsat 8 OLI image based terrestrial water extraction from heterogeneous backgrounds using a reflectance homogenization approach, *Remote Sensing of Environment*. — 171:14–32. DOI: 10.1016/j.rse.2015.10.005 (in Eng.)

CONTENT

A.E. Abetov, A.A. Auyesbek GEOLOGICAL-GEOPHYSICAL CRITERIA OF THE JEZKAZGAN ORE DISTRICT IN CENTRAL KAZAKHSTAN.....	6
K. Akishev, K. Aryngazin, A. Tleulessov, L. Bulyga, V. Stanevich THE USE OF SIMULATION MODELING IN CALCULATING THE PRODUCTIVITY OF THE TECHNOLOGICAL SYSTEM FOR THE PRODUCTION OF BUILDING PRODUCTS WITH FILLERS FROM MAN-MADE WASTE.....	22
V.V. Gerasidi, R.G. Dubrovin, V.V. Kukartsev, T.A. Panfilova, G.V. Stas BOOST SYSTEM DIAGNOSTIC PARAMETERS OF COHERENT GAS PISTON INSTALLATIONS OF MINING ENTERPRISES.....	33
E.M. Elekeev, B.P. Stepanov TO THE ISSUE OF EFFICIENCY OF APPLICATION OF THE SAFETY CONTROL PROGRAM AT NUCLEAR FACILITIES.....	48
A.I. Epikhin, A.A. Stupina, I.A. Panfilov, V.V. Bukhtoyarov, N.A. Shepeta DETERMINANTS FOR ASSESSING THE ENERGY EFFICIENCY OF A COAL MINING ENTERPRISE.....	61
E.A. Efremenkov, E.S. Chavrov, E.P. Khaleyeva, V.V. Tynchenko EVALUATION OF TECHNIQUES FOR DETERMINING THE LOADING OF A CYCLOIDAL SATELLITE ROLLING BEARING.....	72
Zh.A. Zhanabayeva, K.T. Narbayeva, G.K. Ismailova, M.S. Ospanova, J. Rodrigo-Illarri ASSESSMENT OF THE RESERVOIRS IMPACT ON THE MAXIMUM RUNOFF OF THE SYRDARYA RIVER.....	85
M.K. Jexenov, Zh.K. Tukhfatov, E.K. Bektay, R.Sh. Abdinov, G.S. Turysbekova STRUCTURAL-TECHTONIC AND MINERALOGICAL STRUCTURE OF INDER LIFTING FIELD, MINING AND CHEMICAL RAW MATERIALS IN ATYRAU REGION.....	96
G.M. Iskaliyeva^{1,2*}, N.K. Sydyk¹, A.K. Kalybayeva^{1,3}, M.S. Sagat¹, A. Samat USE OF WATER INDICES FOR WATERBODIES IN THE ESIL WATER MANAGEMENT BASIN.....	117
Z.M. Kerimbekova, A.A. Tashimova, G.I. Issayev, E.K. Ibragimova, M.Zh. Makhambetov CALCULATION OF ENVIRONMENTAL AND ECONOMIC DAMAGE CAUSED BY CURRENT SYSTEMS OF SOLID WASTE MOVEMENT IN OIL PRODUCTION..	131
M.Zh. Makhambetov, R. Izimova, G.I. Issayev, N.A. Akhmetov, E.K. Ibragimova ECOLOGICAL-GEOLOGICAL ASSESSMENT OF TECHNOGENICALLY DISTURBED TERRITORIES OF OIL FIELDS OF THE ATYRAU REGION.....	143

A.D. Mekhtiyev, V.V. Yugay, Y.G. Neshina, N.B. Kaliaskarov, P. Madi FIBER-OPTIC SYSTEM FOR CONTROLLING OPEN PIT SIDE ROCK DISPLACEMENT.....	157
A.S. Mussina, G.U. Baitasheva, B.S. Zakirov, Ye.P. Gorbulichева CONDITIONS FOR PREPARING THE SURFACE OF CONTACT PARTS FOR WETTIBILITY WITH MERCURY.....	168
A.N. Muta, R.B. Baimakhan, G.I. Salgarayeva, N. Kurmanbekkyzy, A. Tileikhan STUDY OF THE STRENGTH PROPERTIES OF SOILS COMPOSING THE GEOLOGICAL STRUCTURE OF THE KOK TOBE MOUNTAIN.....	185
B. Mukhambetov, B. Nasiyev, Zh. Kadasheva, R. Abdinov, R. Meranzova ADAPTABILITY OF KOCHIA PROSTRATA (L.) SCHRAD AND CAMPHOROSMA MONSPELIACA AGRICULTURAL ECOSYSTEMS ON SALINE LANDS OF THE NORTHERN CASPIAN DESERT.....	197
M. Nurpeisova, G. Dzhangulova, O. Kurmanbaev, Z. Sarsembekova, A. Ormanbekova, Y.Kh. Kakimzhanov CREATION OF GEODETIC REFERENCE NETWORK FOR MONITORING TRANSPORT INTERCHANGES IN SEISMIC AREAS.....	209
B.T. Ratov, V.L. Khomenko, A.E. Kuttybayev, K.S. Togizov, Z.G. Utepov INNOVATIVE DRILL BIT TO IMPROVE THE EFFICIENCY OF DRILLING OPERATIONS AT URANIUM DEPOSITS IN KAZAKHSTAN.....	224
A.A.Tashimova, G.I. Issayev, Z.M. Kerimbekova, E.K. Ibragimova, A.M. Bostanova ANALYSIS OF THE RESOURCE POTENTIAL OF SOLID OIL PRODUCTION WASTE.....	237
M.K. Tynykulov, O.D. Shoykin, S.O. Kosanov, M.B. Khusainov, Zh.G. Ibraybekov BIOREMEDIATION OF ZONAL SOILS IN AYIRTAY DISTRICT OF NORTH KAZAKHSTAN REGION.....	246
A.S. Urazaliyev, D.A. Shoganbekova, R. Shults, M.S. Kozhakhmetov, G.M. Iskaliyeva INVESTIGATION OF LSMSA APPROACH IN LOCAL GEOID MODELING.....	261

Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the originality detection service Cross Check <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)

ISSN 2518-1483 (Online), ISSN 2224-5227 (Print)

<http://geolog-technical.kz/en/archive/>

Подписано в печать 15.08.2024.

Формат 60x88¹/₈. Бумага офсетная. Печать - ризограф.

15,0 п.л. Тираж 300. Заказ 4.