

ISSN 2518-170X (Online)

ISSN 2224-5278 (Print)

**NEWS OF THE NATIONAL ACADEMY
OF SCIENCES OF THE REPUBLIC
OF KAZAKHSTAN, SERIES OF
GEOLOGY AND TECHNICAL SCIENCES**

№2

2026

ISSN 2518-170X (Online)

ISSN 2224-5278 (Print)



N E W S
OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN,
SERIES OF GEOLOGY AND TECHNICAL
SCIENCES

2 (476)
MARCH – APRIL 2026

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR

ALMATY, 2026

The scientific journal News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences has been indexed in the international abstract and citation database Scopus since 2016 and demonstrates stable bibliometric performance.

The journal is also included in the Emerging Sources Citation Index (ESCI) of the Web of Science platform (Clarivate Analytics, since 2018).

Indexing in ESCI confirms the journal's compliance with international standards of scientific peer review and editorial ethics and is considered by Clarivate Analytics as part of the evaluation process for potential inclusion in the Science Citation Index Expanded (SCIE), Social Sciences Citation Index (SSCI), and Arts & Humanities Citation Index (AHCI).

Indexing in Scopus and Web of Science ensures high international visibility of publications, promotes citation growth, and reflects the editorial board's commitment to publishing relevant, original, and scientifically significant research in the fields of geology and technical sciences.

«Қазақстан Республикасы Ұлттық ғылым академиясының Хабарлары. Геология және техникалық ғылымдар сериясы» ғылыми журналы 2016 жылдан бастап халықаралық реферативтік және ғылымиметриялық Scopus дерекқорында индекстеледі және тұрақты библиометриялық көрсеткіштерді көрсетіп келеді.

Сонымен қатар журнал Web of Science платформасының (Clarivate Analytics, 2018) халықаралық реферативтік және наукометриялық дерекқоры Emerging Sources Citation Index (ESCI) тізіміне енгізілген.

ESCI дерекқорында индекстелуі журналдың халықаралық ғылыми рецензиялау талаптары мен редакциялық этика стандарттарына сәйкестігін растайды, сондай-ақ Clarivate Analytics компаниясы тарапынан басылмды Science Citation Index Expanded (SCIE), Social Sciences Citation Index (SSCI) және Arts & Humanities Citation Index (AHCI) дерекқорларына енгізу қарастырылуда.

Scopus және Web of Science дерекқорларында индекстелуі жарияланымдардың халықаралық деңгейде жоғары сұранысқа ие болуын қамтамасыз етеді, олардың дәйексөз алу көрсеткіштерінің артуына ықпал етеді және редакциялық алқаның геология мен техникалық ғылымдар саласындағы өзекті, бірегей және ғылыми тұрғыдан маңызды зерттеулерді жариялауға ұмтылысын айқындайды.

Научный журнал «News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences» с 2016 года индексируется в международной реферативной и наукометрической базе данных Scopus и демонстрирует стабильные библиометрические показатели.

Журнал также включён в международную реферативную и наукометрическую базу данных Emerging Sources Citation Index (ESCI) платформы Web of Science (Clarivate Analytics, 2018).

Индексирование в ESCI подтверждает соответствие журнала международным стандартам научного рецензирования и редакционной этики, а также рассматривается компанией Clarivate Analytics в рамках дальнейшего включения издания в Science Citation Index Expanded (SCIE), Social Sciences Citation Index (SSCI) и Arts & Humanities Citation Index (AHCI).

Индексирование в Scopus и Web of Science обеспечивает высокую международную востребованность публикаций, способствует росту цитируемости и подтверждает стремление редакционной коллегии публиковать актуальные, оригинальные и научно значимые исследования в области геологии и технических наук.

EDITOR-IN-CHIEF

ZHURINOV Murat Zhurinovich, Doctor of Chemical Sciences, Professor, Academician of IAAS and NAS RK, General Director of the Research Institute of Petroleum Refining and Petrochemicals (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>; <https://www.webofscience.com/wos/author/record/2017489>

DEPUTY EDITOR-IN-CHIEF

ABSADYKOV Bakhyt Narikbayevich, Doctor of Technical Sciences, Professor, Academician of NAS RK, Satbayev University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=6504694468>; <https://www.webofscience.com/wos/author/record/2411827>

EDITORIAL BOARD:

ABSAMETOV Malis Kudysovich, Doctor of Geological and Mineralogical Sciences, Professor, Academician of NAS RK, Director of the U.M. Akhmedsafin Institute of Hydrogeology and Geocology (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=56955769200>; <https://www.webofscience.com/wos/author/record/1937883>

ZHOLTAEV Geroy Zholtaevich, Doctor of Geological and Mineralogical Sciences, Professor, Honorary Academician of NAS RK (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57112610200>; <https://www.webofscience.com/wos/author/record/1939201>

SNOW Daniel, PhD, Associate Professor, Director, Aquatic Sciences Laboratory, University of Nebraska (Nebraska, USA), <https://www.scopus.com/authid/detail.uri?authorId=7103259215>; <https://www.webofscience.com/wos/author/record/1429613>

SELTMANN Reimar, PhD, Head of Petrology and Mineral Deposits Research in the Earth Sciences Department, Natural History Museum (London, Great Britain), <https://www.scopus.com/authid/detail.uri?authorId=55883084800>; <https://www.webofscience.com/wos/author/record/1048681>

PANFILOV Mikhail Borisovich, Doctor of Technical Sciences, Professor at the University of Nancy (Nancy, France), <https://www.scopus.com/authid/detail.uri?authorId=7003436752>; <https://www.webofscience.com/wos/author/record/1230499>

SHEN Ping, PhD, Deputy Director of the Mining Geology Committee of the Chinese Geological Society, Member of the American Association of Economic Geologists (Beijing, China), <https://www.scopus.com/authid/detail.uri?authorId=57202873965>; <https://www.webofscience.com/wos/author/record/1753209>

FISCHER Axel, PhD, Associate Professor, Technical University of Dresden (Dresden, Germany), <https://www.scopus.com/authid/detail.uri?authorId=35738572100>; <https://www.webofscience.com/wos/author/record/2085986>

AGABEKOV Vladimir Enokovich, Doctor of Chemical Sciences, Academician of NAS of Belarus, Honorary Director of the Institute of Chemistry of New Materials (Minsk, Belarus), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>

CATALIN Stefan, PhD, Associate Professor, Technical University of Dresden (Dresden, Germany), <https://www.scopus.com/authid/detail.uri?authorId=35203904500>; <https://www.webofscience.com/wos/author/record/1309251>

JAY Sagin, PhD, Associate Professor, Nazarbayev University (Astana, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57204467637>; <https://www.webofscience.com/wos/author/record/907886>

FRATTINI Paolo, PhD, Associate Professor, University of Milano - Bicocca (Milan, Italy), <https://www.scopus.com/authid/detail.uri?authorId=56538922400>

NURPEISOVA Marzhan Baysanovna, Doctor of Technical Sciences, Professor of Satbayev University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=57202218883>; <https://www.webofscience.com/wos/author/record/AAD-1173-2019>

RATOV Boranbay Tovbasarovich, Doctor of Technical Sciences, Professor, Head of the Department of Geophysics and Seismology, Satbayev University (Almaty, Kazakhstan), <https://www.scopus.com/authid/detail.uri?authorId=55927684100>; <https://www.webofscience.com/wos/author/record/1993614>

RONNY Berndtsson, Professor, Director of the Centre for Advanced Middle Eastern Studies, Lund University (Lund, Sweden), <https://www.scopus.com/authid/detail.uri?authorId=7005388716>; <https://www.webofscience.com/wos/author/record/1324908>

MIRLAS Vladimir, PhD, Professor, Eastern R&D Center, Ariel University (Ariel, Israel), <https://www.scopus.com/authid/detail.uri?authorId=8610969300>; <https://www.webofscience.com/wos/author/record/53680261>

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: «Central Asian Academic Research Center» LLP (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Communications of the Republic of Kazakhstan № KZ50VPY00121155, issued on 05.06.2025
Thematic scope: *geology, hydrogeology, geography, mining and chemical technologies of oil, gas and metals*
Periodicity: 6 times a year.

<http://www.geology-technical.kz/index.php/en/>

© «Central Asian Academic Research Center» LLP, 2026.

БАС РЕДАКТОР

ЖУРЫНОВ Мұрат Жұрынулы, химия ғылымдарының докторы, профессор, ХҒАҚ және ҚР ҰҒА академигі, Мұнай өңдеу және мұнай-химиясы ғылыми-зерттеу институтының бас директоры (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>; <https://www.webofscience.com/wos/author/record/2017489>

БАС РЕДАКТОРДЫҢ ОРЫНБАСАРЫ:

АБСАДЫҚОВ Бақыт Нәрікбайұлы, техника ғылымдарының докторы, профессор, ҚР ҰҒА академигі, Қ.И. Сәтбаев атындағы Қазақ ұлттық техникалық зерттеу университеті (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=6504694468>; <https://www.webofscience.com/wos/author/record/2411827>

РЕДАКЦИЯ АЛҚАСЫ:

ӨБСӘМЕТОВ Мәліс Құдысұлы, геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА академигі, У.М. Ахмедсафин атындағы Гидрогеология және геоэкология институтының директоры (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=56955769200>; <https://www.webofscience.com/wos/author/record/1937883>

ЖОЛТАЕВ Герой Жолтайұлы, геология-минералогия ғылымдарының докторы, профессор, ҚР ҰҒА құрметті академигі (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=57112610200>; <https://www.webofscience.com/wos/author/record/1939201>

СНОУ Дэниел, PhD, қауымдастырылған профессор, Небраска университетінің Су ғылымдары зертханасының директоры (Небраска, АҚШ), <https://www.scopus.com/authid/detail.uri?authorId=7103259215>; <https://www.webofscience.com/wos/author/record/1429613>

ЗЕЛЪТМАНН Раймар, PhD, Жер туралы ғылымдар бөлімінің петрология және пайдалы қазбалар кен орындары саласындағы зерттеулерінің жетекшісі, Табиғи тарих мұражайы (Лондон, Ұлыбритания), <https://www.scopus.com/authid/detail.uri?authorId=55883084800>; <https://www.webofscience.com/wos/author/record/1048681>

ПАНФИЛОВ Михаил Борисович, техника ғылымдарының докторы, Нанси университетінің профессоры (Нанси, Франция), <https://www.scopus.com/authid/detail.uri?authorId=7003436752>; <https://www.webofscience.com/wos/author/record/1230499>

ШЕН Пин, PhD, Қытай геологиялық қоғамының Тау-кен геологиясы комитеті директорының орынбасары, Американдық экономикалық геологтар қауымдастығының мүшесі (Бейжің, Қытай), <https://www.scopus.com/authid/detail.uri?authorId=57202873965>; <https://www.webofscience.com/wos/author/record/1753209>

ФИШЕР Аксель, PhD, қауымдастырылған профессор, Дрезден техникалық университеті (Дрезден, Германия), <https://www.scopus.com/authid/detail.uri?authorId=35738572100>; <https://www.webofscience.com/wos/author/record/2085986>

АГАБЕКОВ Владимир Енокович, химия ғылымдарының докторы, Беларусь ҰҒА академигі, Жаңа материалдар химиясы институтының құрметті директоры (Минск, Беларусь), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>

КАТАЛИН Стефан, PhD, қауымдастырылған профессор, Дрезден Техникалық университеті (Дрезден, Германия), <https://www.scopus.com/authid/detail.uri?authorId=35203904500>; <https://www.webofscience.com/wos/author/record/1309251>

САҒЫНТАЕВ Жанай, PhD, қауымдастырылған профессор, Назарбаев университеті (Астана, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=57204467637>; <https://www.webofscience.com/wos/author/record/907886>

ФРАТТИНИ Паоло, PhD, қауымдастырылған профессор, Бикокок Милан университеті (Милан, Италия), <https://www.scopus.com/authid/detail.uri?authorId=56538922400>

НҮРПЕЙСОВА Маржан Байсанқызы, техника ғылымдарының докторы, Қ.И. Сәтбаев атындағы Қазақ ұлттық техникалық зерттеу университетінің профессоры (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=57202218883>; <https://www.webofscience.com/wos/author/record/AAD-1173-2019>

РАТОВ Боранбай Товбасарович, техника ғылымдарының докторы, профессор, «Геофизика және сейсмология» кафедрасының меңгерушісі, Қ.И. Сәтбаев атындағы Қазақ ұлттық техникалық зерттеу университеті (Алматы, Қазақстан), <https://www.scopus.com/authid/detail.uri?authorId=55927684100>; <https://www.webofscience.com/wos/author/record/1993614>

РОННИ Бердтссон, профессор, Таяу Шығысты заманауи зерттеу орталығының директоры, Лунд университеті (Лунд, Швеция), <https://www.scopus.com/authid/detail.uri?authorId=7005388716>; <https://www.webofscience.com/wos/author/record/1324908>

МИРІАС Владимир, PhD, профессор, Ариэль университетінің Шығыс ғылыми-зерттеу орталығы (Ариэль, Израиль), <https://www.scopus.com/authid/detail.uri?authorId=8610969300>; <https://www.webofscience.com/wos/author/record/53680261>

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктеуші: «Орталық Азия академиялық ғылыми орталығы» ЖШС (Алматы қ.).

Қазақстан Республикасының Ақпарат және коммуникациялар министрлігінің Ақпарат комитетінде 05.06.2025 ж. берілген № KZ50VPY00121155 мерзімдік басылым тіркеуіне қойылу туралы куәлік. Тақырыптық бағыты: *геология, гидрогеология, география, тау-кен ісі, мұнай, газ және металдардың химиялық технологиялары*

Мерзімділігі: жылына 6 рет.

<http://www.geolog-technical.kz/index.php/en/>

ГЛАВНЫЙ РЕДАКТОР

ЖУРИНОВ Мурат Журинович, доктор химических наук, профессор, академик МАН и НАН РК, Генеральный директор НИИ нефтепереработки и нефтехимии (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6602177960>; <https://www.webofscience.com/wos/author/record/2017489>

ЗАМЕСТИТЕЛЬ ГЛАВНОГО РЕДАКТОРА

АБСАДЫКОВ Бахыт Нарикбаевич, доктор технических наук, профессор, академик НАН РК, Казахский национальный исследовательский технический университет им. К.И. Сатпаева (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=6504694468>; <https://www.webofscience.com/wos/author/record/2411827>

РЕДАКЦИОННАЯ КОЛЛЕГИЯ:

АБСАМЕТОВ Малис Кудысович, доктор геолого-минералогических наук, профессор, академик НАН РК, директор Института гидрогеологии и геоэкологии им. У.М. Ахмедсафина (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=56955769200>; <https://www.webofscience.com/wos/author/record/1937883>

ЖОЛТАЕВ Герой Жолтаевич, доктор геолого-минералогических наук, профессор, почетный академик НАН РК (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57112610200>; <https://www.webofscience.com/wos/author/record/1939201>

СНОУ Дэниел, PhD, ассоциированный профессор, директор Лаборатории водных наук Университета Небраски (Небраска, США), <https://www.scopus.com/authid/detail.uri?authorId=7103259215>; <https://www.webofscience.com/wos/author/record/1429613>

ЗЕЛЬТМАНН Раймар, PhD, руководитель исследований в области петрологии и месторождений полезных ископаемых в Отделе наук о Земле Музея естественной истории (Лондон, Великобритания), <https://www.scopus.com/authid/detail.uri?authorId=55883084800>; <https://www.webofscience.com/wos/author/record/1048681>

ПАНФИЛОВ Михаил Борисович, доктор технических наук, профессор Университета Нанси (Нанси, Франция), <https://www.scopus.com/authid/detail.uri?authorId=7003436752>; <https://www.webofscience.com/wos/author/record/1230499>

ШЕН Пин, PhD, заместитель директора Комитета по горной геологии Китайского геологического общества, член Американской ассоциации экономических геологов (Пекин, Китай), <https://www.scopus.com/authid/detail.uri?authorId=57202873965>; <https://www.webofscience.com/wos/author/record/1753209>

ФИШЕР Аксель, PhD, ассоциированный профессор, Технический университет Дрезден (Дрезден, Берлин), <https://www.scopus.com/authid/detail.uri?authorId=35738572100>; <https://www.webofscience.com/wos/author/record/2085986>

АГАБЕКОВ Владимир Еноквич, доктор химических наук, академик НАН Беларуси, почетный директор Института химии новых материалов (Минск, Беларусь), <https://www.scopus.com/authid/detail.uri?authorId=7004624845>

КАТАЛИН Стефан, PhD, ассоциированный профессор, Технический университет Дрезден (Дрезден, Германия), <https://www.scopus.com/authid/detail.uri?authorId=35203904500>; <https://www.webofscience.com/wos/author/record/1309251>

САГИНТАЕВ Жанай, PhD, ассоциированный профессор, Назарбаев университет (Астана, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57204467637>; <https://www.webofscience.com/wos/author/record/907886>

ФРАТТИНИ Паоло, PhD, ассоциированный профессор, Миланский университет Бикокк (Милан, Италия), <https://www.scopus.com/authid/detail.uri?authorId=56538922400>

НУРПЕИСОВА Маржан Байсановна, доктор технических наук, профессор Казахского национального исследовательского технического университета им. К.И. Сатпаева (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=57202218883>; <https://www.webofscience.com/wos/author/record/AAD-1173-2019>

РАТОВ Боранбай Товбасарович, доктор технических наук, профессор, заведующий кафедрой «Геофизика и сейсмология», Казахский национальный исследовательский технический университет им. К.И. Сатпаева (Алматы, Казахстан), <https://www.scopus.com/authid/detail.uri?authorId=55927684100>; <https://www.webofscience.com/wos/author/record/1993614>

РОННИ Бердтссон, профессор, Директор Центра современных ближневосточных исследований, Лундский университет (Лунд, Швеция), <https://www.scopus.com/authid/detail.uri?authorId=7005388716>; <https://www.webofscience.com/wos/author/record/1324908>

МИРЛАС Владимир, PhD, профессор, Восточный научно-исследовательский центр, Университет Ариэля (Ариэль, Израиль), <https://www.scopus.com/authid/detail.uri?authorId=8610969300>; <https://www.webofscience.com/wos/author/record/53680261>

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: ТОО «Центрально-Азиатский академический научный центр» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и коммуникаций и Республики Казахстан № KZ50VPY00121155, выданное 05.06.2025 г.

Тематическая направленность: *геология, гидрогеология, география, горное дело и химические технологии нефти, газа и металлов*

Периодичность: 6 раз в год.

<http://www.geolog-technical.kz/index.php/en/>

CONTENTS

Abetov A.E., Seitzhanov A.K., Samenov Y.R., Zhassymbek S.A. Machine learning and neural networks for lithofacies mapping and reservoir property evaluation of core samples.....	8
Aitkaliyev A.K., Pak Y.N., Kolmakov Y.V., Pak D.Y., Matishev T.V. Application of the X-Ray radiometric method to improve the operational quality control of copper ores during geological testing.....	26
Altay Y.A., Karymsakova N.T., Fedorov A.V., Absadykov D.B., Ismagul G.K. Predictive control of diagnostically significant parameters of acoustic emission using machine learning: new data.....	41
Astanakulov K.D., Kodirova M.G., Saydumarov B., Nersesyan D.V., Kurashkin S.O. Resource potential of ecological and geological systems of allocated soil masses.....	58
Avazov Sh.B., Yodgorov Sh.I., Mukhammadkulov N.M., Aktamov B.U., Khayriddinov B.B. Instrumental seismometric assessment of the impact of technogenic seismic vibrations on the structural stability of historical objects.....	74
Baskanbayeva D., Kalmaganbetov S., Sabirova L., Zholdybayeva G., Koptleuov K. CFD analysis of screw compressor rotor geometry influence on gas compression efficiency in geology.....	92
Borotov A.N., Karshiev F.U., Igamberdiyev A.A., Grigoryan A.G., Sarigo N.V. Ecological aspects of resource conservation in the development of mineral deposits.....	107
Gerasidi V.V., Lisachenko A.V., Apatenko A.S., Sevryugina N.S., Fomin A.Y. Analysis of the application of non-destructive testing methods for assessing the technical condition of mining equipment motors based on vibration parameters.....	121
Hajiyeva A.Z., Ibrahimova L.P., Humbatova Sh.Y., Jafarova F.M. Ecological processes of vertical landscape transformation and their manifestation in the natural landscapes of Nakhchivan.....	135
Jakiyayev D.K., Nussipali R.K.*, Malibekov A.K., Jakiyayev B.D., Zhashen S.Zh. Statistical model of multi-cycle fatigue of steel structural elements under a complex non-uniform stress state.....	149
Jumatov Y.K., Tashbulatov S.B., Pushanov A.N., Kozenkova G.L., Yarygina I.V. Technological substantiation of sustainable mining reclamation of disturbed lands in open pit operations.....	169
Kainenova T.S., Kosmbaeva G.T., Bissembayeva K.T., Otarbayeva A.T., Orazbek A.A. Effectiveness of measures to regulate the development process at the Alibekmola field.....	185

Koishiyeva G.Zh., Yerimova A.Zh., Kurbaniyazov S.K., Agaidarova K.H., Abdraimova K.T.	
Radiological properties and stratigraphic significance of glauconite-bearing formations: implications for paleoenvironments and geochemical zonation.....	203
Kondratiev V.V., Bryukhanova N.N., Kononenko R.V., Gladkikh V.A., Matasova I.Y.	
Environmental and technological assessment of beneficiation tailings reprocessing based on mechanical activation and non-ferrous metals leaching.....	223
Kuldeyev Ye., Negmatov S., Nurpeissova M., Kirgizbaeva D., Donenbayeva N.	
Geodynamic Polygon of the Zhilandy Ore Field: Results and Prospects.....	237
Madibekov A., Christian Opp., Karimov A., Ismukhanova L., Zhadi A.	
Regularities of distribution and accumulation factors of heavy metals in soils of the Zhetysu region (Kazakhstan).....	253
Malikov G.I., Musayeva S.A., Mammadov K.S., Jabiyeva U.M., Jafarova K.V.	
Effects of hydrocarbon fuel quality fluctuations on energy generation efficiency, economic viability, and environmental footprint.....	272
Narbaev M.T., Ismailova G.K., Narbaeva K.T., Burlibayeva D.M., Groll M.	
Assessment of ecological and water management parameters of the Syrdarya River.....	290
Nurmagambetov A.	
On seismic hazard of Almaty and tectonic faults in the city.....	309
Permana A.P., Kasim M., Hutagalung R., Uno D.A.N., Sandi I.N.	
petrographic, provenance, and tectonic setting of Dolokapa formation sandstone, North Gorontalo.....	319
Ratov B.T., Khomenko V.L., Yavorska O.O., Koroviaka Ye.A., Akhmetova N.S.	
Analytical and structural justification of a wellhead sealing device for large-diameter drilling operations.....	335
Skrobala V.M., Popovych V.V., Kopylov V.P., Popovych N.P.	
Comprehensive assessment of heavy metal pollution of a river within a large city.....	355
Sunakbaeva D.Kh., Yuldashbek D.Kh., Aitekova K., Seitbayev K.Zh.	
Monitoring heavy metals in soils and waters of the City of Turkestan: geoecological aspect.....	371
Zakirov M.M., Agzamova I.A., Ochilov G.E., Gulyamov G.D., Xudayberdiyev T.M.	
Analysis of the results of changes in the gas and isotopic composition of groundwater during the period of earthquake preparation.....	385
Zhanbatyrov A.A., Tukhfatov Zh.K., Bektay E.K., Turisbekova G.S., Zhanbatyrov N.A.	
Current state of potash ore development in the Caspian lowlands of the Republic of Kazakhstan: problems and solutions.....	309

NEWS OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC
OF KAZAKHSTAN, SERIES OF GEOLOGY AND TECHNICAL SCIENCES
ISSN 2224-5278
Volume 2.
Number 476 (2026), 399–418

<https://doi.org/10.32014/2026.2518-170X.634>

UDC: 551.1/4
IRSTI: 38.35.21

©Zhanbatyrov A.A.^{*1}, Tukhfatov Zh.K.², Bektay E.K.³, Turisbekova G.S.³,
Zhanbatyrov N.A.¹, 2026.

¹LLC "Tackop Inter", Astana, Kazakhstan;

²NGDU "Dossormunaigas", JSC "Embamunaigas", Atyrau, Kazakhstan;

³Satbayev University, Almaty, Kazakhstan.

E-mail: zaaz1953@mail.ru

CURRENT STATE OF POTASH ORE DEVELOPMENT IN THE CASPIAN LOWLANDS OF THE REPUBLIC OF KAZAKHSTAN: PROBLEMS AND SOLUTIONS

Zhanbatyrov Arun — Candidate of Technical Sciences, LLP Tackop Inter, Astana, Kazakhstan,
E-mail: zaaz1953@mail.ru, <https://orcid.org/0009-0002-8845-931x>;

Tukhfatov Zhasulan — Candidate of Geological and Mineralogical Sciences, NGDU
"Dossormunaigas", JSC "Embamunaigas", Atyrau, Kazakhstan,
E-mail: zh.tuhfatov@kmgep.kz, <https://orcid.org/0009-0007-3301-9997>;

Bektay Erkin — Candidate of Technical Sciences, Satbayev University, Almaty, Kazakhstan,
E-mail: erbekta@gmail.com, <https://orcid.org/0000-0002-5304-5021>;

Turusbekova Gauhar — Candidate of Technical Sciences, Satbayev University, Almaty, Kazakhstan,
E-mail: ipgauharta@gmail.com, <https://orcid.org/0000-0003-1085-6470>;

Zhanbatyrov Nurlan — Master's Student, LLP Tackop Inter, Astana, Kazakhstan,
E-mail: nuriksam@mail.ru, <https://orcid.org/0009-0002-1565-1802>.

Abstract. The article analyzes the current state of preparation for the development of potash and potash-bearing ores in the Pre-Caspian Depression. In 2023, a 25-year license (with the possibility of extension) for the extraction of mineral resources was obtained for the Satimola deposit. "Taskor Inter" LLP developed and approved a Mining Operations Plan for the underground extraction of 25 million tons of potash and potash-bearing ores per year.

Results. Using the Satimola structure as an example, the central site has been selected as the primary object for development, which in turn makes it possible to obtain relevant and comprehensive geological information, ensure degassing of the rock mass, and significantly reduce drilling costs.

For the underground development of potash deposits, it is necessary to train qualified specialists at both higher and intermediate levels - from the study of

mineral raw materials to the justification of technological parameters for extraction and processing (Baryakh A.A et al, 2022). It is proposed to organize such training at the S. Utebayev Atyrau University of Oil and Gas (Zhanbatyrov A.A et al, 2024).

Scientific novelty: The proposed innovative method for preparing a solid mineral deposit for industrial development by underground mining has been implemented at the Satimola structure within the project “Mining Operations Plan for Ore Extraction at the Satimola Deposit.”

Practical value. The authors propose conducting additional studies of the mining-geological, mining-technical, hydrogeological, geotechnical conditions, as well as gas-dynamic phenomena, using modern research methods and scientific instruments.

Keywords: boron-bearing structures, boron-potash salts, carnallite-kieserite rocks, boron minerals - hydroboracite, calborite, boracite.

For citations: Zhanbatyrov A.A., Tuhfatov Zh.K., Bektay E.K., Turisbekova G.S., Zhanbatyrov N.A. *Current State of Potash Ore Development in the Caspian Lowlands of the Republic of Kazakhstan: Problems and Solutions. News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences.* 2026. No.2. Pp. 399–418. DOI: <https://doi.org/10.32014/2026.2518-170X.634>

©Жанбатыров А.А.*¹, Тұхфатов Ж.К.², Бектай Е.Қ.³, Тұрысбекова Г.С.³,
Жанбатыров Н.А.¹, 2026.

¹«Таскор Интер» ЖШС, Астана, Қазақстан;

²«Ембімұнайгаз» АҚ, «Доссормұнайгаз» МГӨБ, Атырау, Қазақстан;

³Сәтбаев Университеті, Алматы, Қазақстан.

E-mail: zaaz1953@mail.ru

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ КАСПИЙ МАҢЫ ОЙПАТЫНДАҒЫ КАЛИЙ КЕНДЕРІН ИГЕРУДІҢ ҚАЗІРГІ ЖАҒДАЙЫ: МӘСЕЛЕЛЕРІ МЕН ШЕШУ ЖОЛДАРЫ

Жанбатыров Арун — техника ғылымдарының кандидаты, «Таскор Интер» ЖШС, Астана, Қазақстан,

E-mail: zaaz1953@mail.ru, <https://orcid.org/0009-0002-8845-931x>;

Тухфатов Жасұлан — геология-минералогия ғылымдарының кандидаты, «Доссормұнайгаз» МГӨБ, «Ембімұнайгаз» АҚ, Атырау, Қазақстан,

E-mail: zh.tuhfatov@kmgcp.kz, <https://orcid.org/0009-0007-3301-9997>;

Бектай Еркін — техника ғылымдарының кандидаты, Сәтбаев университеті, Алматы, Қазақстан,

E-mail: erbekta@gmail.com, <https://orcid.org/0000-0002-5304-5021>;

Тұрысбекова Гаухар — техника ғылымдарының кандидаты, Сәтбаев университеті, Алматы, Қазақстан,

E-mail: ipgauharta@gmail.com, <https://orcid.org/0000-0003-1085-6470>;

Жанбатыров Нурлан — магистрант, "Таскор Интер" ЖШС, Астана, Қазақстан,

E-mail: nuriksam@mail.ru, <https://orcid.org/0009-0002-1565-1802>.

Аннотация. Мақалада Каспий маңы ойпатындағы калий және калий құрамды кендерді игеруге дайындықтың қазіргі жағдайына талдау жасалған. 2023 жылы Сатимола кен орнында 25 жыл мерзімге (ұзарту мүмкіндігімен) пайдалы қазбаларды өндіруге лицензия алынған. «Таскор Inter» ЖШС жерасты тәсілімен жылына 25 млн. тонна калий және калийқұрамды кен өндіруге арналған Тау-кен жұмыстарының жоспарын жасап, келісімнен өткізді.

Борқұрамды кендер мен калий кендерінің маңызды техникалық айырмашылығын атап өткен жөн: су әсерінің қауіптілігі. Қазақстанда тәжірибе тек борқұрамды кендерді өндірумен шектеледі.

Нәтижелері. Сатимола құрылымын мысалға ала отырып, орталық учаске бірінші кезекте игеру нысаны ретінде таңдалған, бұл өз кезегінде өзекті және толық геологиялық ақпарат алуға, тау жыныстары массивін дегазациялауға және бұрғылау шығындарын айтарлықтай төмендетуге және жерасты және жерасты жағдайында жүргізілетін барлау нәтижелері бойынша жаңа деректер алу тиімді игеру мен өнеркәсіптік қауіпсіздікті қамтамасыз етуге мүмкіндік береді.

Калий кен орындарын жерасты тәсілімен игеру үшін минералдық шикізатты зерттеуден бастап өндіру және өңдеу технологиялық параметрлерін негіздеуге дейінгі барлық кезеңдерге қажетті жоғары және орта буын мамандарын дайындау үшін, С.Өтебаев атындағы Атырау мұнай және газ университеті базасында ұйымдастыру ұсынылады.

Ғылыми жаңалығы. Жерасты тәсілімен қатты пайдалы қазбалар кен орнын өнеркәсіптік игеруге дайындаудың ұсынылып отырған инновациялық әдісі Сатимола құрылымында «Сатимола кен орнында рудаларды өндіруге арналған Тау-кен жұмыстарының жоспары» жобасында енгізілген.

Практикалық құндылық. Авторлар кен орнын игерудің тау-геологиялық, тау-техникалық, гидрогеологиялық, геотехникалық жағдайларын және газ-динамикалық құбылыстарын қазіргі заманғы зерттеу әдістері мен ғылыми құралдар негізінде қосымша зерттеуді ұсынады.

Түйін сөздер: тырмалау құрылымдары, бор-калий тұздары, карналлит-кизерит жыныстары, бор минералдары-гидроборацит, калиборит, борацит

©Жанбатыров А.А.^{1*}, Тухфатов Ж.К.², Бектай Е.К.³, Турысбекова Г.С.³,
Жанбатыров Н.А.¹, 2026.

¹ТОО «Taskor Inter», Астана, Казахстан;

²НГДУ «Доссормунайгаз» АО «Эмбаунайгаз», Атырау, Казахстан;

³Satbayev University, Алматы, Казахстан.

E-mail: zaaz1953@mail.ru

СОВРЕМЕННОЕ СОСТОЯНИЕ ОСВОЕНИЯ КАЛИЙНЫХ РУД ПРИКАСПИЙСКОЙ НИЗМЕННОСТИ РЕСПУБЛИКИ КАЗАХСТАН: ПРОБЛЕМЫ И РЕШЕНИЯ

Жанбатыров Арун — кандидат технических наук, ТОО «Taskor Inter», Астана, Казахстан,
E-mail: zaaz1953@mail.ru, <https://orcid.org/0009-0002-8845-931x>;

Тухфатов Жасулан — кандидат геолого-минералогических наук, НГДУ «Доссормунайгаз»,
АО «Эмбаунайгаз», Атырау, Казахстан,
E-mail: zh.tuhfatov@kmgerp.kz, <https://orcid.org/0009-0007-3301-9997>;

Бектай Еркін — кандидат технических наук, Satbayev University, Алматы, Казахстан,
E-mail: erbekta@gmail.com, <https://orcid.org/0000-0002-5304-5021>;

Турусбекова Гаухар — кандидат технических наук, Satbayev University, Алматы, Казахстан,
E-mail: ipgauhartas@gmail.com, <https://orcid.org/0000-0003-1085-6470>;

Жанбатыров Нурлан — магистрант, ТОО «Taskor Inter», Астана, Казахстан,
E-mail: nuriksam@mail.ru, <https://orcid.org/0009-0002-1565-1802>.

Аннотация. *Актуальность.* В статье проведён анализ современного состояния готовности к освоению калийных и калийсодержащих руд Прикаспийской впадины. В 2023 году получена лицензия на добычу полезных ископаемых на месторождении Сатимола сроком на 25 лет с возможностью продления. ТОО «Taskor Inter» разработало и согласовало План горных работ по подземной добыче 25 млн тонн калийных и калийсодержащих руд в год. Следует отметить важное технологическое различие между борсодержащими и калийными рудами, связанное с повышенной опасностью воздействия воды. В Казахстане практический опыт ограничивается преимущественно добычей борсодержащих руд. *Результаты.* На примере структуры Сатимола показано, что центральный участок выбран в качестве приоритетного объекта разработки, что позволяет получать актуальную и полную геологическую информацию, осуществлять дегазацию массива горных пород, существенно снижать затраты на бурение, а также получать новые данные в процессе подземной разведки. Это обеспечивает повышение эффективности разработки и уровня промышленной безопасности. Для освоения калийных месторождений подземным способом рекомендуется создание образовательной и научной базы на базе Атырауский университет нефти и газа имени С. Утебаева для подготовки специалистов высшего и среднего звена, необходимых на всех этапах - от изучения минерального сырья до обоснования технологических параметров добычи и переработки. *Научная*

новизна. Предлагаемый инновационный подход к подготовке месторождения твёрдых полезных ископаемых к подземной разработке реализован в структуре Сатимола в рамках проекта «План горных работ по добыче руд на месторождении Сатимола». *Практическая ценность.* Авторами обоснована необходимость дальнейшего комплексного изучения горно-геологических, горно-технических, гидрогеологических, геотехнических условий и газодинамических процессов разработки месторождения с применением современных методов исследований и научных инструментов.

Ключевые слова: борносные структуры, борно-калийные соли, карналлит-кизеритовые породы, борные минералы - гидроборатит, калиборит, борацит

Introduction. More than 100 years ago, Kazakhstan began and successfully develops hydrocarbon raw materials within the Caspian Basin. The Republic of Kazakhstan has no practical experience in underground mining of potash and potash-containing ores. The current state of readiness for the development of potash and potash-containing ores in the Caspian basin raises reasonable concerns, which may negatively affect the implementation of projects to create and develop the potash industry in the Republic of Kazakhstan in the future. The results of this approach can be seen in Uzbekistan and Turkmenistan during the development of the Garlyk and Tyubegatan fields. The world literature contains materials on the study of the Earth's potassium basins: - The Nepean deposit of potash salts and the Lena-Kirenga potassium-bearing basin of Precambrian age. - Starobinskoye deposit of the Verkhnedevonsky potassium-bearing formation. - Petrikovskoye potassium-bearing deposit of the Upper Devonian age. - The Saskatchewan potassium deposit of the Middle Devonian age. - Cane Creek potash deposit (USA) of medium-carboniferous age. - Verkhnekamskoye deposit of potash salts of the Kungursky stage of the Lower Permian - Verkhnepechorsky basin of potash salts of the Kungurian stage of the Lower Permian, - The Caspian potassium-bearing basin of the Kungurian stage of the Lower Permian. - The Stasfurt potassium-bearing basin of the Upper Permian (Zechstein) age. - The Central European Tsechstein circulatory basin of the Permian age. - Zechstein of Poland of the Late Permian age of potassium salts. - The Kaliningrad (Nivenskoye) potassium deposit of the Permian period. - The Delaware potassium-bearing basin of Late Permian age. - Yorkshire potassium deposit of Late Permian age. - The Central Asian potash basin is confined to the Upper Jurassic and Lower Cretaceous halogen formations. - Industrial accumulations of Paleogene-age potash salts in northern Spain. - The Upper Rhine potassium-bearing basin of tertiary age. - The Precarpathian potassium-bearing basin of Miocene age.

Object of the study. Of the above-mentioned potassium-bearing basins, only in the Caspian salt basin of the Kungurian stage of the lower Permian large-scale industrial deposits of boron-potassium salts unknown to early science were

discovered. The length of the basin is 1000 km, the width is 550 km, and the area is 600 thousand km². Figure 1 shows a forecast map of borates and potash salts in the Caspian basin. The "Central potassium-boron province" has been identified in the potassium-bearing basin. Its length is 400 km, its width is 250 km (Zhanbatyrov et al, 2025).




Figure 1. Forecast map of borates and potash salts of the Caspian lowland. Symbols: 1-contours of domes based on gravimetric and seismic work; 2-stratoisohypses of the salt litter; 3-potassium structures; 4-rake structures; 5 - salt structures of the zone promising for boron and potassium; 6-contours of the central potassium-Boron province.

The object of the study were: 1. Indersky salt massif, located in the southern part of the Central potassium-boron province. The vault area of the structure is 250 km², the salt layer lies at a depth of 55 meters. (Tukhfatov et al, 2024)

Among boron-potassium and potash salts, there are silvinites, silvin-carnallite, carnallite silvinites, kieserite, boracite-halite-kieserite, carnallite-kieserite, carnallite, bischofite-carnallite, bischofite-halite, preobrazhenskite-bischofite-halite and bischofite rocks.

2. The Satimolinsky salt massif, located closer to the center of the Central potassium-boron province of the Caspian Basin. The vault area of the structure is 270 km² and the gypsum layer lies at a depth of 240-250 meters. Potassium and boron-potassium salts of the Satimol structure are composed of polyhalite silvinites, kainite, kieserite-kainite, kainite-polyhalite-carnallite, carnallite-kieserite, kieserite-carnallite rocks with boracite, preobrazhenskite, caliborite, caliborite-preobrazhenskite, preobrazhenskite-caliborite and preobrazhenskite-sulfoborite-caliborite rocks.

3. The Satimolinsky salt massif, located closer to the center of the Central

potassium-boron province of the Caspian Basin. The vault area of the structure is 270 km² and the gypsum layer lies at a depth of 240-250 meters.

Boron-potash salts of the Chelkara salt massif are characterized by complex lithological compositions. Productive breeds are mostly polymineral. Boracite, calborite, and preobrazhenskite are primarily used.

Main content. The study of potash and boron-potash salts has been carried out for several decades at the Atyrau University of Oil and Gas named after Safi Utebayev under the supervision of Academician of the National Academy of Sciences of the Republic of Kazakhstan M.Diarov (Diarov, 2019; Diarov et al, 2016).

The actual chemical salt compositions of the Chelkarsky, Satimolinsky, and Inder structures were analyzed. Since the ore strata have now undergone significant changes in lithological and chemical compositions and it is impossible to draw generalizing conclusions from them, it was necessary to find the gross (based on mass analyses) chemical salt compositions of boron-potash salts of Chelkarsky, Satimolinsky and Inder deposits of the boric province, and first of all, the dependence of individual salt components based on the average composition was clarified. boron-bearing rocks and are shown in Tables 1-3.

Table 1. Average gross chemical composition of boron-bearing horizons of salt massifs (for rocks containing more than 0.5% boron oxide).

Deposits	Samples	Average chemical composition of the components, %											
		B ₂ O ₃	K	Mg	Ca	Na	Cl	SO ₄	CO ₃	Br	H.O.	H ₂ O	The sum of the salts
Chelkar	222	4,34	3,0	2,93	2,69	22,54	40,19	9,37	-	0,102	4,45	10,44	100
Satimola	388	5,13	4,13	3,47	3,14	20,69	35,85	14,03	2,68	0,022	5,17	5,7	100
Inder	999	3,47	5,63	3,40	2,18	23,48	41,83	12,18	0,54	0,039	1,07	6,72	100

Table 2. Salt composition of borosolic ores of boron deposits.

No, p/p	Deposits	Salt compositions (in weight %)							
		B ₂ O ₃	MgCl ₂	MgSO ₄	KCl	CaSO ₄	NaCl	H.O.	MgCO ₃
1	Chelkar	4,34	3,53	3,66	5,72	9,14	57,43	4,45	-
2	Satimola	5,13	0,27	8,15	7,87	10,67	52,6	5,17	3,77
3	Inder	3,47	0,7	4,82	10,73	7,4	59,69	1,07	0,78

Table 3. The actual ratio of gross salt components (in weight. %) boron-potash salts.

Deposits	$\frac{CaSO_4}{B_2O_3}$	$\frac{(MgSO_4 + MgCl_2)}{B_2O_3}$	$\frac{CaSO_4}{MgSO_4 + MgCl_2}$
Chelkar	2,11	1,63	1,27
Satimola	2,08	1,64	1,27
Inder	2,14	1,60	1,34

Despite the complex, diverse lithological, mineralogical and chemical composition of the rocks and the significant remoteness of individual parts of the Central potassium-boron Province, the concentration of boron oxide is strictly related to magnesium and calcium salts. Specifically, the limit of the dependence of boron oxide and magnesium and calcium salts is theoretically determined by the ratio of molecular weights, which was established by academician M. Diarov, based on numerous studies. This approach has received the status of a scientific discovery and is one of the important search features (Tukhfatov et al, 2025; Trapeznikov, 2021). The multicomponent ores of the Caspian lowland require a special approach and a comprehensive study of mineral raw materials, taking into account mining, geological, mining, hydrogeological, geotechnical conditions and gas dynamic phenomena (Asylkhankyzy et al, 2024). These factors are discussed in the monograph "Mining and chemical raw materials for the production of potash and potassium-containing fertilizers" (Zhanbatyrov et al, 2024; Diarov, 2019). Let us briefly consider each factor separately using the example of the structure of Satimol.

Geological conditions. All structures of this type are classified as deposits with difficult mining and geological conditions. The Satimol structure is located in the Central part of the Caspian syncline. Tectonically, the Caspian syncline is characterized by the presence of three structural floors: subsalt, saline and suprasalt (Kasenov et al, 2023). A joint analysis of the relief, the thickness of the sulfate rocks and the morphology and lithology of the salt rocks themselves allowed us to identify four anticlinal structures within the arched part of the Satimola salt massif: the Central, Southeastern, Western, and Kaldybai (Fig. 2). The axes of the anticlinal folds are elongated parallel to the axes of the dome and undulate along the strike.

The central anticlinal structure. The Central anticlinal structure is the most promising for the discovery of new deposits of boron-potassium salts. The most powerful complex of potash and boron-potassium salts is found here. The structure is stretched from northwest to southeast for a distance of 10 km with a width of 2-3 km. The ratio of the long and short axes is 1:3.5. The fold is tilted to the northeast. Along the stretch, the structure sinks to a depth, the capacity of boron-potassium salts is abnormally high. This may be due to the partial preservation of the hinge part of the fold. The potash deposit of the central site was explored in 1967-1973 together with boron-potash salts and eluvial borates. Potash salts are characterized by unstressed morphology and variable mineralogical and chemical composition. The length of the strata is up to 6000m. The capacity ranges from 2m to 35m, with even more in some parts of the field.

The strata of potash and boron-potash salts of the productive zone do not have a rectilinear direction in occurrence, but are complicated by isoclinal folds of various orders, with almost directed axial planes, with hinges sinking in the direction of the general fall of the anticlinal structure. In accordance with this, the outcrops of the layers of the productive zone onto the salt mirror have a complex pattern reflecting the internal tectonics of the salt stratum.



Figure 2. Diagram of the tectonic structure of the dome of the salt dome of Satimola.
Scale 1:50000.

1-anticlinal structures; 2-areas of predominantly monoclinic occurrence of salt rocks.

Southeastern anticlinal structure It stretches from northeast to southwest. The length is up to 6 km. The ratio of the long and short axes is 1:3. Adjacent synclinal structures, elongated in a northwesterly direction, are located everywhere on the dome next to the identified anticlinal structures. Their core consists of younger age salt deposits. Judging by the geological situation of the dome and the data of oriented samples taken from exploratory wells, the angles of rock fall are mostly steep, reaching $50-60^\circ$. However, there are areas where potash and boron salts lie in a relatively flat (no steeper than at an angle of $30-45^\circ$) monocline.

Kaldybai anticlinal structure. It is strongly elongated in a northeasterly direction. The ratio of the axes is at least 1:7. Boron-potash and potash salts of the southwest wing were discovered by wells №№2, 34, 20, 29, 23 and 380. The fold is isoclinal in nature with steep angles of incidence of the wings. According to the data of the oriented sample of well No. 34, the angle of incidence of the southwest wing of the fold is 75° . The northeastern wing of the Kaldybai structure is also composed of potash and boron-potash salts discovered by wells No.93, 10, 4, 22 and 382. In the section of well No. 4 (range 606-610 m), the boron-bearing pack is

represented by brownish-red rock salt, boracite-anhydrite rocks. The upper boron-bearing pack in well No. 23 is based on silvinites, and the lower part is weakly boron-bearing. The boron horizon was uncovered by well No. 23 at a depth of 593-599 m and is lithologically represented by boracite-carnallite-halite-kieserite rock with a weighted average content of B₂O₃ - 3.60% and K₂O - 1.99%. Well No. 34 uncovered the upper boron bearing pack at a depth of 384-397 m under the silvinitic formation. It consists of two boron-bearing formations bearing boracite mineralization, separated by a layer of rock salt. The upper stratum is composed of rock salt with a weighted average content of B₂O₃ - 0.78%, the lower one is composed of halite-kieserite rock with a content of B₂O₃ - 4.35%.

Western anticlinal structure. The structure is elongated from northwest to southeast. Judging by the data of structural prospecting wells and kernoscopy, the outcrops to the salt mirror of the rocks of the productive zone in the southwestern wing repeat the character of the outer wing of this site in their configurations. Apparently, the tectonic forces that displaced the dome wing simultaneously affected the nature of the occurrence of boron salts in the Western anticline. The length of the structure is 17 km. A peculiar complex of boron-potassium, potash, and potassium-magnesium salts of various compositions has been discovered in the salt column, confined to certain horizons in the middle part of the stratigraphic salt section.

The studied potash deposits are a series of irregular morphologically stratified and lenticular bodies. They have been opened by many wells. The potassium occurrences are cut by single wells.

The outlet of the layers of the potassium-bearing zone to the salt mirror has a complex configuration due to the peculiarities of the internal tectonics of the salt massif and is shown in Fig.3.

Thus, the mineralization of the Central potash-boron province of the Caspian Basin has no analogues in the world practice in terms of quality, quantity and variety of boron raw materials. Boron-potassium salts containing boron, potassium and magnesium are easily enriched complex mineral raw materials. In addition, the presence of rare earth metals significantly increases the potential value of the mineral resources of the Caspian Basin. (Freyman, 2020; Asylkhankyzy et al., 2020)

The presence of lithium, rubidium, and caesium was found in all lithological varieties of salt rocks in the dome. The upper horizons of the salt domes of the Caspian basin have been washed away by groundwater, and huge amounts of salts have been removed from them. When salts are removed, insoluble minerals accumulate above the salt mirror in the form of a kind of tire, called a "plaster hat" or cap. The main part of precious metals and REM are concentrated in insoluble salt sediment and represent practically natural storages. The content of lithium, rubidium and caesium in boron concentrates of four technological samples of salts of the Satimola dome is shown in Table 4.

Table 4. The content of lithium, rubidium and caesium in boron concentrates of four technological samples of salts of the Satimola dome.

No. p/p	Samples	Mineral composition	The content of rare and scattered elements, in %		
			Li	Rb	Cs
1	Technological sample No. 1. Boron salts (sle.No.512, int.803-818m)	Halite, polyhalite, caliborite, clay	0,0066	0,0030	0,0002
2	Technological sample concentrate No. 1	Caliborite, polyhalite, clay	0,0140	0,005	0,0006
3	Technological test No. 17. Boron salts. (square No.513, int.452,5-456m)	Halite, sylvin, polyhalite, caliborite, anhydrite	0,0072	H.o.	0,0002
4	Technological sample concentrate No. 17	Preobrazhenskite, polyhalite,, anhydrite, clay	0,0225	H.o.	H.o.
5	Technological test No. 18. Boron salts	Halite, sylvin, polyhalite, caliborite, anhydrite	0,0066	0,0020	H.o.
6	Technological sample concentrate No. 18	Preobrazhenskite, polygalite, caliborite, clay	0,01	0,0030	H.o.
7	Technological test No. 19. Boron ores. (square No. 514, int.1080-1086m).	Halite, polyhalite, caliborite, clay	H.o.	H.o.	0,0001
8	Technological sample concentrate No. 19.	Caliborite, polyhalite, clay	H.o.	H.o.	0,0001

Mining conditions. The mining conditions of the Satimol structure are considered in detail in the Mining Plan. (Plan gornyxh rabot, 2023) Taking into account the morphology of the productive layers of the Satimola deposit, the following development systems are planned:

- ore mining using combine harvesters (flat and inclined formations);
- drilling and blasting method of ore extraction (inclined and steeply falling formations);

The discovery of reserves of potash and boron salts of the Satimola deposit is carried out in stages.

– the first stage is the construction of mine shafts No. 1 and No. 2, which open the Central section of the deposit.

– the second stage is the opening of the South flank of the deposit with shafts No. 4 and No.5.

– the third stage is the construction of ventilation shaft No. 3 with a diameter of 8 m on the North-West flank of the deposit, which provides the necessary amount of fresh air to ventilate the mine workings with a maximum production of 25.0 million tons of potash ore per year.

Hydrogeological conditions. The central potassium-boron province is located within the North Caspian groundwater basin. During the exploration of the Satimola deposit in 1979-1985 and with additional research at the mine shaft site in 2008, 61 hydrogeological wells were drilled and pilot filtration work was carried out.

The groundwater of the Satimol structure is connected to the suprasalt hydrogeological floor, the average capacity of which reaches 360 m. Table 5 shows the rocks composing the hydrogeological section.




Figure 3. Geological and lithological map of the salt mirror of the salt dome structure of Satimol. Compiled by K.Kamashev, 1979.

1-sulfate stratum (gypsum and anhydrites); 2-eluvial borates (hydroboracite, ulexite), upper halite zone; 3-rock salt; 4-potassium salts (sylvinites, carnallites, polyhalites); 5-anhydrites; 6-rock salt with borate inclusions (caliborite, asharite, hydroboracite) productive boron bearing zone; 7-rock salt; 8-potash salts (sylvinites, carnallites, polyhalites, kainites); 9-boron-bearing rocks (boracite, preobrazhenskite, caliborite, asharite, hydroboracite).

Table 5. Rocks forming the hydrogeological section.

Age (index)	Breeds	Depth, m	Filtration coefficient, m/day
Qm _h v	Lenses of sand in clays	10	0,07
Qb-Qn _h z	Lenses of sand in clays	10-60	2,52
N ₂ ³ ap - N ₂ ³ ak	Lenses of sand in clays	60-90	10 ⁻⁶
N ₂ ³ ap - N ₂ ³ ak	Lenses of sand in clays	90-130	2,26
N ₂ ³ ap - N ₂ ³ ak	Lenses of sand in clays	130-190	10 ⁻⁷
K ₂	Clays, chalk, marl, seashell	190-210	1,0
K ₁	Clay	210-230	10 ⁻⁶
J _{1,2}	Clay	230-280	10 ⁻⁶
T	Clay	230-360	10 ⁻⁶
P ₁ k ^p	Gypsum, clay, borate ores	230-360	0,0077

The generalization of all hydrogeological materials on the Satimola deposit allows us to draw the following conclusions on the assessment of water flows into the mine workings during the development of potash salts.

1) The expected inflow of groundwater into the mine shafts during their sinking will not exceed 35 dm³/s (126m³/h) – from the flooded stratum of the Upper Pliocene Akchagyl horizon (N23ak). The sinking of mine shafts should provide for reliable isolation of aquifers, followed by their overlap with structural elements of the mine shaft.

2) The only probable external source of groundwater inflows into the mining operations carried out for the development of salts may be exploration geological wells cut by mining operations. These wells will work as through filters. In reality, flows from them are possible only from the weakly water-bearing Lower Permian Kungur sulfate zone (P1kp), composed of relatively stable rocks capable of ensuring the safety of that part of the exploration wells that uncovered these rocks. Due to the low filtration parameters of these rocks, inflows from such wells will not exceed 1-2 liters/s for any period of rock salt development. However, in order to avoid cutting exploration geological wells by mining operations and ensure the safety of mining operations, it is recommended to leave well sights near.

3) The salt column is practically anhydrous. The source of water supply to the mine workings (subject to the safety of the ceiling, borehole pillars and the tightness of the mine shafts) may be rare intra-formation leaching cavities with a volume not exceeding 20-30 m³, with a maximum daily inflow of up to 20 m³. In this regard, given the insignificance of brine occurrence, there is no need for special measures, since it is possible to limit the special equipment available in the underground mine, only providing for the availability of storage tanks, followed by the involvement of brines in the drilling process.

Geotechnical conditions. Engineering and geological studies on the structure of Satimol were carried out at all stages of geological exploration, as a result of which

representative data for the deposit were obtained. In addition, to substantiate the design decisions, the experience of work at the Inderskaya exploration mine was used, based on the general genesis of salt rocks, their almost similar composition (the Caspian salt basin) and the uniformity of structures (salt domes). (Kasenov et al, 2024; Kasenov et al, 2024). Research at the Satimola deposit was carried out by VNIIG (1967) and GIGHS (1979 - 1985). From the entire complex of engineering and geological studies, the following initial data were selected: for the suprasalt stratum - the results of the GIGHS, for the salt stratum - the work of 2008-2010, as the most representative for the deposit in terms of area, depth, and number of tests.

The GIGHS examined samples taken from nine wells, which fairly fully represent both the Central Site (eluvial borates) and the deposit as a whole. A total of about 300 rock samples were taken.

According to the conclusion of GIGHS, the cover layer of the Satimola deposit is generally characterized as relatively homogeneous in terms of strength.

The sulfate stratum containing boron mineralization is characterized by stable strength parameters, with the exception of massive gypsum anhydrite layers, which are characterized by average values of compressive strength of 496 kg/cm² and tensile strength of 55 kg/cm².

The corresponding figures for gypsum are 237 and 27.3 kg/cm². Gypsum-clay rocks lying directly above the mineral strata are noticeably weaker than the main massif: the compressive strength is 95.4 kg/cm², and the tensile strength is 26.3 kg/cm². For salt rocks, the compressive strength ranges from 233 to 519 kg/cm², and tensile strength ranges from 17 to 21 kg/s.

Based on the results of engineering and geological studies, the GIGHS concluded that there are no large zones (sites) with weakened rock stability, weathering zones, and tectonic fragmentation within the Central Area, which could be identified and shown in the scale of the geological map of the field being presented. Such areas are small and local. Given the fact that the Central Site is the most structurally complex element of the Satimol structure, this conclusion can be confidently extrapolated to the entire deposit.

Gas dynamic phenomena (gas content). The main feature of the mining conditions during the operation of the Satimola deposit is the presence of gas content here. The first signs of the presence of gas phenomena on the structure of Satimol were noted in 1964 during prospecting for boron-potassium salts. From 1965 to 1968, special gas drilling studies were carried out at the structure in 28 structural exploration wells relatively evenly spaced over the area.

Analysis of gas logging materials showed that all recorded gas occurrences are confined to the salt column of the structure and, as a rule, are located at depths exceeding 500 m in 25 wells out of 28; only in three wells weak gas occurrences are recorded at depths of 300-400 m. Moreover, in two of them, these manifestations are confined to the upper horizons of the salt stratum, and in one – to the bottoms of the gypsum-anhydrite stratum and the tops of the salt. Thus, a conclusion was made about the gas content of the salt stratum.

In 1983-84, at the stage of detailed exploration of the Central section of the field (eluvial borates), gas logging was carried out by the LGKS-4AP serial station in 16 wells during drilling and component-by-component analysis of drilling fluid samples for marginal hydrocarbon gases using a gas-capping thermograph of the XG-1G type, which allows determining the content of the air mixture in gases of the maximum hydrocarbons from CH₄ to C₆H₁₄. As a result, a conclusion was made about the gas permeability of kaprock and its low gas content.

1. Only folded dislocations are developed at deposit No. 99, and there are no disjunctive tectonic disturbances. There is one extended intraformational zone associated with the movement of salt masses inside the core of the structure, a fracture zone, a parallel crumpling zone in the host salts, and several size-limited fractured zones. These tectonic elements have no outlet to the salt mirror and are associated with breccia and conglomerate-like textures, especially of carnallite rocks, with somewhat weakened physical and mechanical properties (Tukhfatov et al, 2024).

2. All workings were completed without fastening, isolated collapses were observed in breccated carnallite rock.

3. Strength coefficient of potash salts according to Protodyakonov 1-2, boron salts 3-5. The stability assessment was carried out by measuring the workings with a section of 7.5 square meters. m. The sides of the workings approached each other by 8 mm in 110 days, and the roof by up to 3mm. Thus, it was concluded that the rocks are stable.

4. The only source of brine occurrences are the deaf cavernous cavities of leaching filled with extremely saturated uterine brines.

5. The recommended capacity of the ceiling water-proof rear sight is 100 m.

6. The gas content of the deposit is related to both the host rocks and ore layers. The gas is in a free, bound and micro-connected state. The distribution of free gas is extremely uneven and ranges from 0 to 11.5 m³/m³. The average free gas content in rocks is rock salt, non-explosive - from 0 to 2, with an average of 0.39 m³/m³; explosive - 5.3 m³/ m³, carnallite-kieserite-polyhalite - 0.29 m³/m³. Free gases consist of methane (70-99%), heavy hydrocarbons and nitrogen are present in small amounts (0.6-2.2%).

The distribution of bound gases is uneven from 0.005 to 2.83 m³/t.

Explosive rocks have the highest gas content, which is on average 4 times higher than non-explosive rocks. Composition nitrogen (41-88%), methane (5.7-40.2%), hydrogen (4.08-7.55%), carbon dioxide (0.12-2.67%). In the total gas content (0.48 m³/m³) of bound gases, 1-10%, the rest are free. The gas permeability of salts is very low and does not exceed 9.5x10⁻² m. darcy.

By the nature of the manifestation, they are divided into:

– Usual gas emissions from the surface of mine workings (pressure less than 10 kPa) for rocks with carnallite up to 100x10⁻⁴m³/m²/min, on average 52-574m³/m²/min; silvinites – 51x10⁻⁴m³/m²/min and 32 x10⁻⁴m³/m²/min; rock salt -17 x10⁻⁴m³/m³/min and 14 x10⁻⁴m³/m²/min, respectively.

– Souffle gas emissions occur from boreholes and wells or cracks that have exposed accumulations of free gas. Of the 164 wells analyzed, souffle gas emissions were recorded in 55, and in some souffles were observed 10-16 times, while there was an outflow or discharge of drilling fluid, and at high pressure, drilling tools. The maximum measured pressure was 4 MPa, the highest calculated pressure determined by the fracture factor of the drilling tool was 38 MPa. At a pressure of more than 10 kPa, a rock pin is ejected into the hole, and at a higher pressure, a hand-held drilling tool can be ejected. The density of souffles is 1.3 for every 100 p.m of drilling.

There are two types of sudden emissions: actual sudden emissions and sudden rashes. The latter occur after blasting operations and in all cases are associated with some kind of geological complications, where the rocks are weakened and have little strength. Sudden emissions of rock and gas form spherical and elliptical shapes with an average volume of 16.9 m³.

According to the intensity of emissions from blasting operations, they are divided into five types: microbursts of up to 3 tons/discharge (11.2% of all emissions), low-strength - 3-20 tons/discharge (29.9%), medium-strength - 20-150 tons/discharge (41.1%), high-strength - 150-600 tons/discharge (15.9%) and powerful - more than 600 tons/emission (1.9%). Most of the high-intensity and powerful emissions come from the host rock salt.

It should be noted that in the course of geological exploration in 2008-2010, 24 cases of gas release from wells were recorded at the Satimola field. For the most part, gas occurrences were noted in the form of foaming of drilling mud, and only in three cases was the release of the solution, while the maximum calculated gas pressure was about 9 MPa. The difference between these data and the experience of deposit No. 99 is explained by the depth of the salt relative to the wellhead - 350 m (Satimol) and 0 m (Inder) during underground drilling from the mine and the associated pressure of the column of flushing liquid. With a washing liquid density of about 1.3 g/cm³, the initial pressure at the entrance to the salt layer at a depth of 350 m will be about 4.5 MPa and increases by 1.27 MPa every 100 m. Such an excess of the solution pressure over the gas pressure does not allow the latter to enter the borehole, in addition, the process of colmatation of cracks and pores during drilling generally eliminates any manifestations (of course, at a lower gas pressure).

The methods of forecasting gas occurrences at field No. 99 developed during this period were:

– Core. When the core is divided into discs with a length less than its diameter, a sudden release will occur regardless of the method of penetration. The method is reliable for rocks with a strength of less than 140 kgf/cm², but with higher strength it is not always reliable, since the division into discs does not always occur in this case.

Most of the emissions in boron salts are confined to areas with a maximum content of boron oxide.

– The acoustic forecasting method is based on measuring the noise level of salt when dissolved in water. It is applicable to carnallite rocks in the drilling and blasting method of sinking.

These methods are planned to be applied at the Satimola field during mining operations and, if necessary, supplemented to predict gas events.

Materials and methods. A comprehensive study of potash and boron-potassium salts was carried out using various methods:

- Continuous sampling (core) of boron-potash salts obtained as a result of drilling boreholes.
- Detailed description of the rocks.

- Lithological, mineralogical and chemical study of potash and boron salts by determining mineralogical compositions under a microscope in immersion liquids; conducting complete chemical analyses with determination of B_2O_3 , K, Mg, Ca, Na, Cl, SO_4 , CO_3 and insoluble residue; recalculation by component compositions of rocks into salt compositions.

- Study of the physical and mechanical properties of the mountain range, taking into account the modeling of loads under different conditions and positions.

- Study of gas dynamic phenomena on special stands with core samples, taking into account different porosity, pressure and temperature changes.

Results and discussions. All conditions and phenomena that arise and affect technological processes during mining operations were studied.

1. The complex geological conditions of the structures of the Caspian Lowland and the multicomponence of ores should be noted. Ore bodies are represented by stratified, plate-lenticular and lenticular deposits, while they are often not consistent in terms of thickness and mineral composition. The presence of noble and rare earth metals increases the potential value of the object. The separate location of eluvial borates makes it possible to develop a separate project, regardless of the main part of potash and potash-containing ores.

When preparing deposits of potash and potash-containing ores by the underground method, it is proposed to explore the central part in detail by conducting a complex of exploration, and to explore the remaining areas by conducting advanced operational exploration. (Kasenov et al, 2024; Yue Hao et al, 2025).

The specified geological conditions must be taken into account when developing project documents for mining.

2. The hydrogeological conditions are simple, but given the high dependence of potash ores on the effects of water, it is necessary to plan water protection measures for the entire period of operation of the facility.

3. Mining conditions are also very difficult when developing a deposit underground. Given the high variability of geological conditions, mining of each ore deposit must be approached individually.

4. The geotechnical conditions have been studied, but it is necessary to obtain additional data immediately before developing a specific geological unit in real production conditions.

5. Gas dynamic phenomena, according to research, will occur during the development of horizons below 350 meters.

This research is funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (grant No. AP14870189).

Conclusions. All geological, mineralogical, and gas emission studies were conducted on extracted cores in laboratory conditions at research institutes in the Soviet Union and Kazakhstan, as well as during boron ore mining at the Inder uplift and underground exploration of deposit No. 99.

Pilot industrial development is required for underground mining of potash and potash-bearing ores to clarify technological parameters of the mining systems at this stage (Zhanbatyrov A.A., 2024). These provisions should be incorporated into the current Code of the Republic of Kazakhstan "On Subsoil and Subsoil Use," similar to the approach used in uranium deposit development.

Based on the studies performed, the following conclusions can be drawn:

1. The Mining Plan for the extraction of ores from the Satimola deposit takes into account all the features of mining, geological, mining, hydrogeological, geotechnical conditions and gas dynamic phenomena.

2. The following underground mining systems have been adopted:

- ore mining using combine harvesters (flat and inclined formations);
- drilling and blasting method of ore extraction (inclined and steeply falling strata).

3. Local development planning has been adopted in certain areas with difficult mining and geological conditions.

4. It provides for the use of precious and rare earth metals, i.e. complex processing of mineral raw materials.

5. Provide for the training of mining engineers and technicians for the underground mining of potash deposits at Atyrau University of Oil and Gas.

6. To organize scientific research laboratories for the study of the mineral resources of the Caspian Lowland.

7. To ensure the development of regulatory documents on industrial safety in the underground mining of potash and potash-containing ores.

8. To amend the current Code of the Republic of Kazakhstan "On Subsoil and Subsoil Use", which provides for pilot production to adjust the technological parameters of underground mining systems in order to efficiently and safely develop potash and potash-containing ores.

References

Asylkhankyzy A., Seitmagzimova G.M. (2024) *Peredovye tekhnologii proizvodstva kaliynykh udobreniy iz perspektivnogo syrya* [Advanced Technologies of Potash Fertiliser Production from Promising Raw Materials]. *The Open Chemical Engineering Journal*. – Vol. 18, e18741231330310. (in Russian).

Asylkhankyzy A., Seitmagzimova G.M., Petrov M.M., Dzhanmuldaeva Zh.K. (2020) *Issledovanie sposobov pererabotki kaliynykh rud mestorozhdeniya Chelkar* [Study of processing methods of potash

ores from the Chelkar deposit]. Bulletin of the Ural State Technical University. Kazakhstan-British Technical University, 17(2). – P. 44-52. (in Russian).

Baryakh A.A., Smirnov E.V., et al. (2022) Kaliynaya promyshlennost' Rossii: Problemy ratsional'nogo i bezopasnogo nedropol'zovaniya [Potash Industry of Russia: Problems of Rational and Safe Subsoil Use]. Journal Mining Industry. – No. 1. – P. 41-50. (in Russian).

Diarov M.D. (2019) Fundamentalnye printsipy nakopleniya i formirovaniya promyshlennykh mestorozhdeniy bora sovместno s kaliynymi solyami solenosnogo basseyna [Fundamental principles of accumulation and formation of industrial boron deposits together with potash salts of the salt basin]. Atyrau. – P. 33. (in Russian)

Diarov M.D. (2019) Svidetelstvo o vnesenii v Gosudarstvennyy reestr obektov avtorskogo prava No. 5375 ot 19.09.2019. Treugolnoe pravilo [Certificate of entry into the State Register of Copyrighted Objects. – No. 5375, dated 19.09.2019. The Triangle Rule. (in Russian).

Diarov M.D., Saburov O.E. (2016) Geologiya galogennykh boratov i kaliynykh soley Prikaspiyskoy vpadiny [Geology of halogen borates and potash salts of the Pre-Caspian Depression]. Mon, Izd-vo "Evero", Almaty, TOO "Color Media". – P. 238-240. (in Russian).

Freyman G.G. (2020) Otchet s podschetom zapasov kaliynykh soley po Zhilyanskomu mestorozhdeniyu v Aktyubinskoy oblasti po sostoyaniyu na 01.01.2014 g. [Report with reserve estimation of potash salts at the Zhilyansk deposit, Aktobe region]. Almaty. (in Russian)

Kasenov T.I., Zhanbatyrov A.A. (2023) Etapy osvoeniya mineral'nykh resursov mestorozhdeniya Satimola [Stages of development of mineral resources of the Satimola deposit]. Bulletin "Atyrau oil and gas university named by S.Utebayev", Atyrau. – P. 4-11. (in Russian)

Kasenov T.I., Zhanbatyrov A.A. (2024) K voprosu o vybore sposoba razrabotki mestorozhdeniya Satimola [On the selection of the mining method for the Satimola deposit]. Bulletin "Atyrau oil and gas university named by S.Utebayev". – No.3(71). – P. 41-44. (in Russian)

Kasenov T.I., Zhanbatyrov A.A., Telibaev B.K. (2023) Plan zavochnykh rabot na mestorozhdenii Satimola [Verification work plan at the Satimola deposit]. Astana, Ltd, 2024. – No. 88-ML (November 22). – P. 49. (in Russian)

Kasenov T.I., Zhanbatyrov N.A., Telibaev B.K., Seitmukanov M.S. (2024) Zayavka na patent RK «Sposob podgotovki mestorozhdeniya tverdykh poleznykh iskopaemykh k osvoeniyu podzemnym sposobom» [Patent application of the Republic of Kazakhstan "Method for preparing a solid mineral deposit for underground development"]. IP «KenArAl», Astana. – P. 69. (in Russian)

Plan gornykh rabot po dobyche rud mestorozhdeniya Satimola (2023) [Mining operations plan for ore extraction at the Satimola deposit]. Book 1 in 2 parts. Astana, Taskor Inter LLP. (in Russian)

Trapeznikov D. (2021) Geologicheskoe stroenie Inderskogo solyanogo kupola [Geological structure of the Inder salt dome]. Mining Echo. – No.3. – P. 17-22. (in Russian)

Tukhfatov Zh.K., Dzheksenov M.K., Bektai Y.K., Abdinov R.Sh., Turisbekova G.S. (2024) Strukturno-tektonicheskoe i mineralogicheskoe stroenie otlozheniy Inderskogo podnyatiya [Structural-tectonic and mineralogical structure of the Inder uplift deposits]. Izvestiya NAN RK, seriya geologicheskikh issledovaniy. – No. 4(465). – P. 96-116. (in Russian)

Tukhfatov Zh.K., Dzheksenov M.K., Bektai Y.K., Turisbekova G.S., Shiderin B.N. (2025) Poiskovye raboty na gorno-khimicheskoe syrye v odnom iz solyanykh kupolov Prikaspiyskoy vpadiny [Prospecting for mining-chemical raw materials in one of the salt domes of the Caspian depression]. News of the National Academy of Sciences of the Republic of Kazakhstan, Series of Geology and Technical Sciences. – No. 3(471). – P. 252-267. (in English).

Yue Hao, Minghui Yu, Wei Weijie, Shengli Yang, Jinwang Zhang, Zhaohui Wang, Zewei Wang, Yukun Liu, Junrui Ge. (2025) Obzor metodov i oborudovaniya dlya dobychi kaliynykh udobreniy [A Review of Potash Mining Methods and Equipment]. Mining, Metallurgy & Exploration. – Vol. 42. – No. 6. – P. 3515-3545. (in English).

Zhanbatyrov A.A. (2024) Upravlenie investitsiyami pri poetapnom osvoenii mestorozhdeniy poleznykh iskopaemykh [Investment management in phased development of mineral deposits]. Bulletin "Atyrau oil and gas university named by S.Utebayev". – No. 3(71). – P. 141-145. (in Russian)

Zhanbatyrov A.A., Iskakov R.M. (2024) Sozdanie nauchno-issledovatel'skikh tsentrov na

sovmestnoy baze predpriyatiy i universitetov [Creation of research centers based on cooperation between enterprises and universities]. Bulletin "Atyrau oil and gas university named by S.Utebayev". – No. 3(71). – P. 150-153. (in Russian)

Zhanbatyrov A.A., Kasenov T.I., Iskakov R.M., Telibaev B.K., Zhanbatyrov N.A. (2024) Gornokhimicheskoe syrye dlya proizvodstva kaliynykh i kaliysoderzhashchikh udobreniy [Mining and chemical raw materials for the production of potash and potassium-containing fertilizers]. Astana. – P. 369. (in Russian)

Zhanbatyrov A.A., Zhanbatyrov N.A. (2025) Sovremennoe sostoyanie razvitiya kaliynoy otrasli v RK [Current state of development of the potash industry in the Republic of Kazakhstan]. Bulletin "Atyrau oil and gas university named by S.Utebayev". – No. 4. (in Russian).

Publication Ethics and Publication Malpractice in the journals of the Central Asian Academic Research Center LLP

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the journals of the Central Asian Academic Research Center LLP implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The Central Asian Academic Research Center LLP follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/ or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the Central Asian Academic Research Center LLP.

The Editorial Board of the Central Asian Academic Research Center LLP will monitor and safeguard publishing ethics.

Requirements for articles design for publication in the journal are available on the websites:

**[www:nauka-nanrk.kz](http://www.nauka-nanrk.kz)
<http://www.geolog-technical.kz/index.php/en/>
ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)**

Managing Editor: *T. Apendiev*
Editors: *D.S. Alenov, A.Shormakova*
Computer layout: *G.D. Zhadyranova*

Signed for print: April 10, 2026
Format: 70×90 1/16. 26.5 printed sheets. Order No. 2.