

ISSN 2518-170X (Online),  
ISSN 2224-5278 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ  
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ

# Х А Б А Р Л А Р Ы

---

---

## ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК  
РЕСПУБЛИКИ КАЗАХСТАН

## NEWS

OF THE ACADEMY OF SCIENCES  
OF THE REPUBLIC OF KAZAKHSTAN

ГЕОЛОГИЯ ЖӘНЕ ТЕХНИКАЛЫҚ ҒЫЛЫМДАР  
СЕРИЯСЫ



СЕРИЯ  
ГЕОЛОГИИ И ТЕХНИЧЕСКИХ НАУК



SERIES  
OF GEOLOGY AND TECHNICAL SCIENCES

### 3 (423)

МАМЫР – МАУСЫМ 2017 ж.  
МАЙ – ИЮНЬ 2017 г.  
MAY – JUNE 2017

ЖУРНАЛ 1940 ЖЫЛДАН ШЫҒА БАСТАҒАН  
ЖУРНАЛ ИЗДАЕТСЯ С 1940 г.  
THE JOURNAL WAS FOUNDED IN 1940.

ЖЫЛЫНА 6 РЕТ ШЫҒАДЫ  
ВЫХОДИТ 6 РАЗ В ГОД  
PUBLISHED 6 TIMES A YEAR

АЛМАТЫ, ҚР ҰҒА  
АЛМАТЫ, НАН РК  
ALMATY, NAS RK

Б а с р е д а к т о р ы

э. ғ. д., профессор, ҚР ҰҒА корреспондент-мүшесі

**И.К. Бейсембетов**

Бас редакторының орынбасары

**Жолтаев Г.Ж.** проф., геол.-мин. ғ. докторы

Р е д а к ц и я а л қ а с ы:

**Абаканов Т.Д.** проф. (Қазақстан)  
**Абишева З.С.** проф., корр.-мүшесі (Қазақстан)  
**Алиев Т.** проф., академик (Әзірбайжан)  
**Бакиров А.Б.** проф., (Қырғыстан)  
**Беспәев Х.А.** проф. (Қазақстан)  
**Бишимбаев В.К.** проф., академик (Қазақстан)  
**Буктуков Н.С.** проф., корр.-мүшесі (Қазақстан)  
**Бұлат А.Ф.** проф., академик (Украина)  
**Ганиев И.Н.** проф., академик (Тәжікстан)  
**Грэвис Р.М.** проф. (АҚШ)  
**Ерғалиев Г.Х.** проф., академик (Қазақстан)  
**Жуков Н.М.** проф. (Қазақстан)  
**Кенжалиев Б.К.** проф. (Қазақстан)  
**Қожахметов С.М.** проф., академик (Қазақстан)  
**Конторович А.Э.** проф., академик (Ресей)  
**Курскеев А.К.** проф., академик (Қазақстан)  
**Курчавов А.М.** проф., (Ресей)  
**Медеу А.Р.** проф., корр.-мүшесі (Қазақстан)  
**Мұхамеджанов М.А.** проф., корр.-мүшесі (Қазақстан)  
**Нигматова С.А.** проф. (Қазақстан)  
**Оздоев С.М.** проф., академик (Қазақстан)  
**Постолатий В.** проф., академик (Молдова)  
**Ракишев Б.Р.** проф., академик (Қазақстан)  
**Сейтов Н.С.** проф., корр.-мүшесі (Қазақстан)  
**Сейтмуратова Э.Ю.** проф., корр.-мүшесі (Қазақстан)  
**Степанец В.Г.** проф., (Германия)  
**Хамфери Дж.Д.** проф. (АҚШ)  
**Штейнер М.** проф. (Германия)

«ҚР ҰҒА Хабарлары. Геология мен техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде 30.04.2010 ж. берілген №10892-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18, <http://nauka-nanrk.kz/geology-technical.kz>

---

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2017

Редакцияның Қазақстан, 050010, Алматы қ., Қабанбай батыра көш., 69а.

мекенжайы: Қ. И. Сәтбаев атындағы геология ғылымдар институты, 334 бөлме. Тел.: 291-59-38.

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р

д. э. н., профессор, член-корреспондент НАН РК

**И. К. Бейсембетов**

Заместитель главного редактора

**Жолтаев Г.Ж.** проф., доктор геол.-мин. наук

Р е д а к ц и о н н а я к о л л е г и я:

**Абаканов Т.Д.** проф. (Казахстан)  
**Абишева З.С.** проф., чл.-корр. (Казахстан)  
**Алиев Т.** проф., академик (Азербайджан)  
**Бакиров А.Б.** проф., (Кыргызстан)  
**Беспаяев Х.А.** проф. (Казахстан)  
**Бишимбаев В.К.** проф., академик (Казахстан)  
**Буктуков Н.С.** проф., чл.-корр. (Казахстан)  
**Булат А.Ф.** проф., академик (Украина)  
**Ганиев И.Н.** проф., академик (Таджикистан)  
**Грэвис Р.М.** проф. (США)  
**Ергалиев Г.Х.** проф., академик (Казахстан)  
**Жуков Н.М.** проф. (Казахстан)  
**Кенжалиев Б.К.** проф. (Казахстан)  
**Кожаметов С.М.** проф., академик (Казахстан)  
**Конторович А.Э.** проф., академик (Россия)  
**Курскеев А.К.** проф., академик (Казахстан)  
**Курчавов А.М.** проф., (Россия)  
**Медеу А.Р.** проф., чл.-корр. (Казахстан)  
**Мухамеджанов М.А.** проф., чл.-корр. (Казахстан)  
**Нигматова С.А.** проф. (Казахстан)  
**Оздоев С.М.** проф., академик (Казахстан)  
**Постолатий В.** проф., академик (Молдова)  
**Ракишев Б.Р.** проф., академик (Казахстан)  
**Сейтов Н.С.** проф., чл.-корр. (Казахстан)  
**Сейтмуратова Э.Ю.** проф., чл.-корр. (Казахстан)  
**Степанец В.Г.** проф., (Германия)  
**Хамфери Дж.Д.** проф. (США)  
**Штейнер М.** проф. (Германия)

«Известия НАН РК. Серия геологии и технических наук».

**ISSN 2518-170X (Online),**

**ISSN 2224-5278 (Print)**

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10892-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,

<http://nauka-nanrk.kz/geology-technical.kz>

---

© Национальная академия наук Республики Казахстан, 2017

Адрес редакции: Казахстан, 050010, г. Алматы, ул. Кабанбай батыра, 69а.

Институт геологических наук им. К. И. Сатпаева, комната 334. Тел.: 291-59-38.

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of Economics, professor, corresponding member of NAS RK

**I. K. Beisembetov**

Deputy editor in chief

**Zholtayev G.Zh.** prof., dr. geol-min. sc.

E d i t o r i a l b o a r d:

**Abakanov T.D.** prof. (Kazakhstan)  
**Abisheva Z.S.** prof., corr. member. (Kazakhstan)  
**Aliyev T.** prof., academician (Azerbaijan)  
**Bakirov A.B.** prof., (Kyrgyzstan)  
**Bespayev Kh.A.** prof. (Kazakhstan)  
**Bishimbayev V.K.** prof., academician (Kazakhstan)  
**Buktukov N.S.** prof., corr. member. (Kazakhstan)  
**Bulat A.F.** prof., academician (Ukraine)  
**Ganiyev I.N.** prof., academician (Tadjikistan)  
**Gravis R.M.** prof. (USA)  
**Yergaliev G.Kh.** prof., academician (Kazakhstan)  
**Zhukov N.M.** prof. (Kazakhstan)  
**Kenzhaliyev B.K.** prof. (Kazakhstan)  
**Kozhakhmetov S.M.** prof., academician (Kazakhstan)  
**Kontorovich A.Ye.** prof., academician (Russia)  
**Kurskeyev A.K.** prof., academician (Kazakhstan)  
**Kurchavov A.M.** prof., (Russia)  
**Medeu A.R.** prof., corr. member. (Kazakhstan)  
**Muhamedzhanov M.A.** prof., corr. member. (Kazakhstan)  
**Nigmatova S.A.** prof. (Kazakhstan)  
**Ozdoyev S.M.** prof., academician (Kazakhstan)  
**Postolatii V.** prof., academician (Moldova)  
**Rakishev B.R.** prof., academician (Kazakhstan)  
**Seitov N.S.** prof., corr. member. (Kazakhstan)  
**Seitmuratova Ye.U.** prof., corr. member. (Kazakhstan)  
**Stepanets V.G.** prof., (Germany)  
**Humphery G.D.** prof. (USA)  
**Steiner M.** prof. (Germany)

**News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.**

**ISSN 2518-170X (Online),**

**ISSN 2224-5278 (Print)**

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 10892-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,  
<http://nauka-nanrk.kz/geology-technical.kz>

---

© National Academy of Sciences of the Republic of Kazakhstan, 2017

Editorial address: Institute of Geological Sciences named after K.I. Satpayev  
69a, Kabanbai batyr str., of. 334, Almaty, 050010, Kazakhstan, tel.: 291-59-38.

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 3, Number 423 (2017), 228 – 235

UDC 539.3(043.3)

A. Zh. Seytmuratov<sup>1</sup>, D. M. Zharylgapova<sup>1</sup>, N. K. Medeubaev<sup>2</sup>, A. A. Ibraeva<sup>1</sup>

<sup>1</sup>Korkyt Ata Kyzylorda State University, Kyzylorda, Kazakhstan,

<sup>2</sup>E. A. Buketov Karaganda State University, Karaganda, Kazakhstan.

E-mail: angisin\_@mail.ru, djm.06@mail.ru

## APPLIED TASKS OF PLATES FLUCTUATION UNDER MORE DIFFICULT BOUNDARY CONDITIONS

**Abstract.** The results on the research of own and compelled fluctuations of flat elements taking into account lamination of material of plate, rheological viscous properties, anisotropies, etc. are given in this work.

At the research of harmonious waves in deformable bodies, the concept of phase speed as the speed of environment condition change is implemented; at the same time, the phase speed is expressed through the frequencies of own fluctuations and therefore the research of harmonious waves distribution has a direct bearing on the problems of definition of own forms and frequencies of the fluctuations limited in terms of plates.

More difficult fluctuation of rectangular flat element is the fluctuation when two of the opposite edges are hinge-supported, and two other edges have different types of fixing or they are free from tension.

This class of tasks leads to the transcendental equations for determination of frequencies of own fluctuations which can be solved both numerically and analytically.

The transcendental frequency equations can be reduced to algebraic ones and to investigate the influence of both boundary conditions at the edges of rectangular plate or rectangular flat element and parameters of geometrical and mechanical character on the frequencies of own fluctuations of rectangular flat elements.

**Key words:** plates, anisotropy, own fluctuations, compelled fluctuations, transcendental, equations, rheology.

We will consider flat element as isotropic uniform elastic plate of constant thickness [2].

We will be limited to the task solution based on approximate equation of quartic cross fluctuations

$$A_0 \frac{\partial^4 W}{\partial t^4} - A_1 \Delta \frac{\partial^2 W}{\partial t^2} + A_2 \Delta^2 W + \frac{\partial^2 W}{\partial t^2} = \Phi(f_z, f_{jz})(j = x, y) \quad (1)$$

where coefficients  $A_j$  are equal to

$$A_0 = \frac{h^2(7-8\nu)}{12b^2(1-\nu)}; \quad A_1 = \frac{2h^2(2-\nu)}{3(1-\nu)}; \quad A_2 = \frac{2h^2b^2}{3(1-\nu)} \quad (2)$$

$\nu$  – Poisson ratio;  $b$  – speed of cross waves distribution in plate material.

As plate edges ( $y = 0; l_2$ ) are hinge-supported, the solution of the equation (1) we will find in

$$W(x, y, t) = \exp\left(i \frac{b}{h} \xi t\right) \sum_{k=1}^{\infty} W_k(x) \sin\left(\frac{k\pi y}{l_2}\right) \quad (3)$$

Substituting (3) in the equation (1), for  $W_k$  we get ordinary differential equation

$$\frac{d^4 W_k}{dx^4} + B_0 \frac{d^2 W_k}{dx^2} + B_1 W_k = 0 \quad (4)$$

where coefficients  $B_0, B_1$  are equal to

$$B_0 = \left[ \frac{A_1}{A_2} \xi^2 \left( \frac{b}{h} \right)^2 - 2 \left( \frac{k\pi}{l_2} \right)^2 \right];$$

$$B_1 = \left[ \left( \frac{k\pi}{l_2} \right)^4 + \frac{A_0}{A_2} \xi^4 \left( \frac{b}{h} \right)^4 - \frac{A_1}{A_2} \xi^2 \left( \frac{b}{h} \right)^2 \left( \frac{k\pi}{l_2} \right)^2 - \frac{1}{A_2} \left( \frac{b}{h} \right)^2 \xi^2 \right] \quad (5)$$

We will write down the common solution of the equation (4) as

$$W_k(x) = C_1 \left[ \frac{\cos(a_0 x)}{a_0^n} + \frac{\cos(a_1 x)}{a_1^n} \right] + C_2 \left[ \frac{\cos(a_0 x)}{a_0^n} + \frac{\cos(a_1 x)}{a_1^n} \right] +$$

$$+ C_3 \left[ \frac{\sin(a_0 x)}{a_0^m} + \frac{\sin(a_1 x)}{a_1^m} \right] + C_4 \left[ \frac{\sin(a_0 x)}{a_0^m} + \frac{\sin(a_1 x)}{a_1^m} \right], \quad (6)$$

where  $C_j$  - integration constants;  $a_i, a_j$  - roots of the characteristic equation

$$a^4 + B_0 a^2 + B_1 = 0 \quad (7)$$

and are equal to

$$a_{0,1} = \sqrt{\frac{B_0}{2}} \pm \sqrt{\left( \frac{B_0}{2} \right)^2 - B_1} \quad (8)$$

Integers  $(n, m)$  are got out of solution simplification condition at the satisfaction of boundary condition at the left edge  $x = 0$ , and other boundary conditions at  $x = l_1$  lead to the transcendental equation for the determination of frequencies of the plate own fluctuations.

We will analyze transcendental frequency equations of the first point.

In the beginning we will consider the simplest transcendental equation

$$\alpha_0 \cos(\alpha_0 l_1) \sin(\alpha_1 l_1) - \alpha_1 \sin(\alpha_0 l_1) \cos(\alpha_1 l_1) = 0. \quad (9)$$

We will implement designations

$$l = \frac{l_1}{h}; \quad \alpha_{0,1}^1 = \sqrt{\frac{B_0^1}{2}} \pm \sqrt{\left( \frac{B_0^1}{2} \right)^2 - B_1^1}; \quad (10)$$

$$B_0^n = [(2 - \nu)\xi^2 - 2\gamma]; \quad \gamma = \left( \frac{\pi k h}{l_2} \right)^2;$$

$$B_1^i = \left[ \gamma^2 + \frac{7 - 8\nu}{8} \xi^4 - (2 - \nu)\gamma\xi^2 - \frac{3}{2}(1 - \nu)\xi^2 \right]$$

and further we will lower strokes for simplicity.

As sines and cosines from any argument are equal

$$\sin z = \sum_{i=0}^{\infty} (-1)^i \frac{z^{2i+1}}{(2i+1)!}; \quad \cos z = \sum_{j=0}^{\infty} (-1)^j \frac{z^{2j}}{(2j)!};$$

the equation (9) is equivalent to the following

$$\alpha_0 \alpha_1 \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} (-1)^{i+j} \frac{\alpha_1^{2i} \alpha_0^{2j} - \alpha_0^{2i} \alpha_1^{2j}}{(2i+1)!(2j)!} l^{2(i+j)} = 0 \quad (11)$$

If to accept that size is determined from the formula (8) with plus sign under the root, then it follows that this root does not turn zero at any values  $y, \nu, \xi$ .

Therefore, in the beginning it is possible to put  $a_1 = 0$  or for  $\xi$  we get the equation

$$\xi^4 - \frac{8\left[(2-\nu)\gamma + \frac{3}{2}(1-\nu)\right]}{(7-8\nu)}\xi^2 + \frac{8\gamma^2}{(7-8\nu)} = 0; \quad (12)$$

which roots are equal to

$$\begin{aligned} \xi_{1,2} = (7-8\nu)^{\frac{2}{2}} \sqrt{4\left[(2-\nu)\gamma + \frac{3}{2}(1-\nu)\right]} \pm \\ \pm \sqrt{8(1+\nu^2)\gamma^2 + 3\gamma(1-\nu)(2-\nu) + \frac{9}{4}(1-\nu)^2} \end{aligned} \quad (13)$$

as the ranks in the formulae of trigonometrical functions meeting ranks in the equation (11), equivalent to the equation (9) also meeting, at the research of private equation (11) it is possible to be limited to the final number of the first composed.

Having taken first three composed in the ranks (11), we will write down it as

$$\begin{aligned} \alpha_0\alpha_1(\alpha_1^2 - \alpha_0^2) \left\{ \frac{1}{3}l^2 - \frac{1}{30}(\alpha_1^2 + \alpha_0^2)l^4 + \right\} \times \\ \times \left[ \frac{1}{840}(\alpha_1^4 + \alpha_0^2\alpha_1^2 + \alpha_0^4) + \frac{1}{360}\alpha_0^2\alpha_1^2 \right] l^6 + \dots \} = 0 \end{aligned} \quad (14)$$

Roots from the formula  $\alpha_1 = 0$  are equal (13). The value  $(\alpha_1^2 - \alpha_0^2)$  is other than zero at any  $y, \nu, \xi$ .

If in formula (14) to take only two first terms, then we get

$$(\alpha_1^2 + \alpha_0^2) - 10l^{-2} = 0$$

or

$$B_0 - 10l^{-2} = 0 \quad (15)$$

and frequency equation

$$\xi^2 = \frac{2\gamma + 10l^{-2}}{(2-\nu)}; \quad (16)$$

which positive root is equal to

$$\xi = \sqrt{\frac{2\gamma + 10l^{-2}}{(2-\nu)}} \quad (17)$$

If in formula to take all three first terms, then we will get

$$\left[ (\alpha_1^4 + \alpha_0^4) + \frac{10}{3}\alpha_0^2\alpha_1^2 \right] - 28(\alpha_1^2 + \alpha_0^2)l^{-2} + 280l^{-4} = 0 \quad (18)$$

or

$$\left[ B_0^2 + \frac{4}{3}B_1 \right] - 28B_0l^{-2} + 280l^{-4} = 0$$

and the frequency equation corresponding to them

$$\begin{aligned} \left[ (2-\nu)^2 + \frac{7+8\nu}{6} \right] \xi^4 - \left[ (2-\nu) \left( \frac{16}{3}\gamma + 28l^{-2} \right) + 2(1-\nu) \right] \xi^2 + \\ + \left[ \frac{16}{3}\gamma^2 + 56\gamma l^{-2} + 280l^{-4} \right] = 0, \end{aligned} \quad (19)$$

which has two positive roots.

It is similarly possible to take first four and more terms in the formula (11) and to get more exact frequency equation and corresponding frequencies  $\xi$ .

To find the frequency equation from ranks of the equation (11) it is necessary to find out condition of deduction legitimacy of finite number of terms in the ranks (11).

We will apply Dalamber's principle of ranks convergence to the ranks in the equation (11). We will get

$$\left| \frac{\alpha_0^2 \alpha_1^2 l^2}{(2i+3)(2j+2)} \right| \leq q^2 < 1 \quad (20)$$

where  $0 < q < 1$ .

It follows from the inequality (20) that

$$|a_0^2 a_1^2| \leq q_{i,j}^2 = q_{i,j}^2 = q^2 \frac{(2i+3)(2j+2)}{l^2} \quad (21)$$

The analysis of inequality (21) shows that it is correct when performing inequality

$$-\left(\frac{8}{7-8\nu}\right) q_{i,j}^2 \leq \xi^4 - 2D\xi^2 + E \leq \left(\frac{8}{7-8\nu}\right) q_{i,j}^2 = C_{i,j}^2$$

where coefficients  $D, E$  are equal to

$$D = \frac{4 \left[ (2-\nu)\gamma + \frac{3}{2}(1-\nu) \right]}{(7-8\nu)}; \quad E = \frac{8\gamma^2}{(7-8\nu)}$$

or inequalities

$$D^2 - E \leq C_{i,j}^2 \quad (22)$$

At the set parameters of geometrical and mechanical character from the inequality (22) it is possible to define necessary number of the first terms in ranks (11) to find the frequency equation relative frequencies  $\xi$ .

We will consider transcendental equation frequency equation

$$2 - \frac{a_0^2 + a_1^2}{a_0 a_1} \sin(a_0 l_1) \sin(a_1 l_1) - 2 \cos(a_0 l_1) \cos(a_1 l_1) = 0 \quad (23)$$

As well as transcendental equation (9), the equation (23) is equivalent to the following

$$a_0 a_1 \left\{ 2 \left[ 1 - \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} (-1)^{i+j} \frac{a_0^{2i} a_1^{2j}}{(2i)!(2j)!} J^{2(i+j)} \right] - \right. \\ \left. - (a_0^2 + a_1^2) \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} (-1)^{i+j} \frac{a_0^{2i} a_1^{2j}}{(2i+1)!(2j+1)!} J^{2(i+j+1)} \right\} = 0 \quad (24)$$

It follows from the equation, that, first,  $a_1 = 0$  also we got frequencies (13).

We will write down the equation (24), having written out the first terms

$$\left\{ (a_0^2 + a_1^2) J^2 - \frac{1}{6} (5a_0^4 + 5a_1^4 + a_0^2 a_1^2) J^4 + \right\} \times \\ \times \frac{1}{90} [a_0^6 + a_1^6 + 7a_0^2 a_1^2 (a_0^2 a_1^2)] J^6 + \dots \} = 0 \quad (25)$$

From (25) it also follows that it can be supposed  $(a_0^2 + a_1^2) = 0$  and we will get

$$B_0 = 0 \quad \text{or} \quad \xi^2 - \frac{2\gamma}{(2-\nu)} = 0 \quad (26)$$



which root is equal to

$$\xi = \sqrt{\frac{2\gamma}{(2-\nu)}} \quad (27)$$

Similarly, we can put approximately

$$(a_0^2 + a_1^2) - \frac{J^2}{6}(5a_0^4 + 5a_1^4 + a_0^2 a_1^2) = 0 \quad (28)$$

and we will get frequency equation

$$(5B_0^2 - 9B_1) - \frac{6}{J^2} B_0 = 0 \quad (29)$$

having positive roots.

We will consider more difficult transcendental equation

$$4[Q(Q - a_0^2 - a_1^2) + a_0^2 a_1^2][1 - \cos(a_0 l_1) - \cos(a_1 l_1)] + \\ + 2\left[\frac{a_0^3}{a_1^3}(Q - a_1^2)^2 + \frac{a_1^3}{a_0^3}(Q - a_0^2)^2\right] \sin(a_0 l_1) - \sin(a_1 l_1) = 0, \quad (30)$$

where

$$Q = \frac{3-2\nu}{7-4\nu} \left[ \frac{\xi^2}{h^2} - 2 \left( \frac{\pi k}{l_2} \right)^2 \right]$$

which is equivalent to the following

$$2[Q_0(Q_0 - a_0^2 - a_1^2) + a_0^2 a_1^2] \left[ 1 - \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} (-1)^{i+j} \frac{a_0^{2i} a_1^{2j}}{(2i)!(2j)!} J^{2(i+j)} \right] + \\ + [a_0^4(Q_0 - a_1^2)^2 + a_1^4(Q_0 - a_0^2)^2] \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} (-1)^{i+j} \frac{a_0^{2i} a_1^{2j}}{(2i+1)!(2j+1)!} J^{2(i+j+1)} = 0 \quad (31)$$

$$Q_0 = \left( \frac{3-2\nu}{7-4\nu} \right) (\xi^2 - 2\gamma).$$

We will write out the first terms in the equation (31)

$$[(a_0^2 + a_1^2)L_1 + L_2]J^2 - \left[ L_1 \left( \frac{a_0^4 + a_1^4}{12} + \frac{a_0^2 a_1^2}{2} \right) + L_2 \left( \frac{a_0^2 + a_1^2}{6} \right) \right] J^4 + \\ + \left\{ \frac{1}{360} L_1 [a_0^6 + a_1^6 + 15a_0^2 a_1^2 (a_0^2 + a_1^2)] \right\} + \quad (32) \\ + \frac{1}{20} L_2 \left( a_0^4 + a_1^4 + \frac{10}{3} a_0^2 a_1^2 \right) J^6 + \dots = 0;$$

$$L_1 = [Q_0(Q_0 - a_0^2 - a_1^2) + a_0^2 a_1^2];$$

$$L_2 = [a_0^4(Q_0 - a_1^2)^2 + a_1^4(Q_0 - a_0^2)^2].$$

Being limited by the first term in the formula (32), we will get

$$(a_0^2 + a_1^2)L_1 + L_2 = 0 \quad (33)$$

or

$$Q_0^2(B_0^2 - 2B_1) - 2Q_0 B_0 B_1 + 2B_1^2 + B_0 [Q_0(Q_0 - B_0) + B_1] = 0 \quad (34)$$

The frequency equation (34) is the algebraic equation relatively to  $\xi$  and already depends on the parameter  $\gamma$ .

Equations of higher order relatively to  $\xi$  already depend on the parameter  $\gamma$ .

It is also possible to consider other boundary value problems.

Thus, transcendental frequency equations can be reduced to algebraic and to investigate influence of both boundary conditions at the edges of rectangular plate or rectangular flat element, and parameters of geometrical and mechanical character on frequencies of own fluctuations of rectangular flat elements.

We will generalize outcomes for rectangular plate or flat element which material satisfies viscoelastic model of Maxwell.

Let we have a rectangular uniform isotropic plate.

In this case, we will find the solution of approximate quartic equation

$$W = \exp\left(\frac{b}{h}\xi t\right) \sum_{k=1}^{\infty} W_k \sin\left(\frac{pky}{J_2}\right), \quad (35)$$

where  $\xi$  - complex frequency which valid part defines the law of attenuation of fluctuations and imaginary part determines frequencies of own fluctuations.

For  $W_k$  we get ordinary differential equation

$$\frac{d^4 W_k}{dx^4} - \overline{B_0} \frac{d^2 W_k}{dx^2} + \overline{B_1} W_k = 0 \quad (36)$$

where  $\overline{B_0}, \overline{B_1}$  are equal to

$$\begin{aligned} \overline{B_0} &= \left[ 2\gamma + \frac{A_1}{A_2} \frac{b}{h} \left( \frac{b}{h} \xi^2 + \frac{1}{\tau} \xi \right) \right]; \\ \overline{B_1} &= \left[ \frac{A_0}{A_2} \left( \frac{b}{h} \right)^4 \xi^4 + 2 \frac{A_0}{A_2 \tau} \left( \frac{b}{h} \right)^3 \xi^3 + \left( \frac{b}{h} \right)^2 \times \right. \\ &\quad \left. \left( \frac{1}{A_2} + \frac{1}{\tau^2} \frac{A_0}{A_2} + 2\gamma \frac{A_1}{A_2} \right) \xi^2 + \frac{1}{A_2 \tau} \left( \frac{b}{h} \right) \times \left( 1 + 2\gamma \frac{A_1}{A_2} \right) \xi + \gamma^2 \right] \end{aligned} \quad (37)$$

Coefficients  $A_j$  are provided in the previous paragraphs.

We will write down the common solution of the equation (36)

$$\begin{aligned} W_k &= C_1 \left[ \frac{ch(a_0 x)}{a_1^n} + \frac{ch(a_1 x)}{a_1^n} \right] + C_2 \left[ \frac{ch(a_0 x)}{a_0^n} + \frac{ch(a_1 x)}{a_1^n} \right] + \\ &+ \left[ \frac{sh(a_0 x)}{a_0^m} + \frac{sh(a_1 x)}{a_1^m} \right] + C_4 \left[ \frac{sh(a_0 x)}{a_0^m} + \frac{sh(a_1 x)}{a_1^m} \right], \end{aligned} \quad (38)$$

i.e. instead of trigonometrical functions we have hyperbolic ones.

All considered regional tasks leading to the transcendental equations are solved similarly. Transcendental equations are got from previous where values  $a_j$  are necessary to replace with values  $ia_j$ , where  $i$  - imaginary unit.

For example, the transcendental equation (9) passes into the equation

$$a_0 ch(a_0 l_1) sh(a_1 l_1) - a_1 sh(a_0 l_1) ch(a_1 l_1) = 0 \quad (39)$$

which is equivalent to the following

$$a_0 a_1 \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \frac{a_1^{2i} a_1^{2j} a_0^{2i}}{(2i+1)!(2j)!} l^{2(i+j)} = 0;$$

$$a_{0,1} = \sqrt{\frac{B_0}{2}} \pm \sqrt{\frac{B_0^2}{4} - B_1} \quad (40)$$

One of the frequency equations of the formula (40) follows from the condition  $a_j = 0$  and we get

$$\begin{aligned} \xi^4 + \frac{2}{\tau_0} \xi^3 + \frac{8}{(7-8\nu)} \left[ (2-\nu)\gamma + \frac{(7-8\nu)}{8\tau_0^2} + \frac{3}{2}(1-\nu) \right] \xi^2 + \\ + \frac{12(1-\nu)}{(7-8\nu)\tau_0} [1 + 2(2-\nu)\gamma] \xi + \frac{8}{(7-8\nu)} \gamma^2 = 0, \end{aligned} \quad (41)$$

Which coincides with the frequency equation of the approximate equation of quartic cross fluctuations (1) for rectangular hinge-supported plate on all four parties of plate, and it has two complex conjugate roots.

#### REFERENCES

- [1] Filippov I.G. To the nonlinear theory of viscoelastic isotropic environments. Kiev: Applied mechanics, 1983. Vol. 19, N 3. P. 3-8.
- [2] Filippov I.G., Filippov S.I. Equations of fluctuation of piecewise uniform plate of variable thickness. MTT. 1989. N 5. P. 149-157.
- [3] Filippov I.G., Filippov S.I., Kostin V.I. Dynamics of two-dimensional composites. Works of the Int. conference on mechanics and materials, the USA, Los Angeles, 1995. P. 75-79.
- [4] Seytmuratov A.Zh. Passing of shift waves through anisotropic non-uniform and transversal isotropic cylindrical layer. Dep. in KazgostINTI No. 189-B 96. Release p.17. Almaty 1996.
- [5] Seytmuratov A.Zh. Approximate equations of cross fluctuation of plate under the surface. Theses of reports of scientific technical conference "Environmental problem and environmental management". K-Orda, 1996.
- [6] Seytmuratov A.Zh. Specified equations of fluctuation of viscoelastic plate under the surface of deformable environment. Theses of reports of Zhakhayev KPTI scientific and technical conference. K-Orda, 1996.
- [7] Seytmuratov A.Zh. Fluctuations of infinite strip of plate under the surface. Dep. in VINITI No. 3399-B 96 from 22.11.96. M., 1996.
- [8] Filippov I.G., Cheban V.G. Mathematical theory of fluctuations of elastic and viscoelastic plates and cores. Kishinev: Shtiintsa, 1988. P. 190-193.
- [9] Materials of international scientifically-practical conference "The Science: theory and practice". Belgorod, 2005. P. 47-50.
- [10] Seytmuratov A.Zh., Umbetov U. Modeling and forecasting of dynamics of multicomponent deformable environment: Monograph. Taraz, 2014. P. 171-176.
- [11] Seytmuratov A.Zh. Metod of decomposition in the theory of fluctuation of two-layer plate in building constructions. PGS. 2006. N 3. M. P. 31-32.
- [12] Seytmuratov A.Zh. Determination of frequency of own fluctuations of plate. Messenger of KazNU, mathematics, mechanics, computer science series. 2010. N 4(67).

А. Ж. Сейтмұратов<sup>1</sup>, Д. М. Жарылғапова<sup>1</sup>, Н. К. Медеубаев<sup>2</sup>, А. А. Ибраева<sup>1</sup>

<sup>1</sup>Қорқыт Ата атындағы Қызылорда мемлекеттік университеті, Қазақстан,

<sup>2</sup>Бөкетов атындағы Қарағанды мемлекеттік университеті, Қазақстан

#### АСА КҮРДЕЛІ ШЕКАРАЛЫҚ ШАРТТАР КЕЗІНДЕГІ ПЛАСТИНАЛАР ТЕРБЕЛІСІНІҢ ҚОЛДАНБАЛЫ ЕСЕБІ

**Аннотация.** Берілген жұмыста пластинкадан құралған материалдың қатпарлылығын, реологиялық тұтқыр қасиеттерін, анизотропиясын және т.б. зерттеу нәтижелері есере отырып шешілген меншікті және еріксіз тербелістер есебін қарастырылған.

Зерттеу нәтижесінде гармоникалық толқындардың деформацияланатын денелер жағдайындағы фазалық жылдамдығын, орта күйінің өзгеру жылдамдығы деп қарастырады, бұл ретте фазалық жылдамдық жиілігі меншікті тербеліс арқылы өрнектеледі, сондықтан гармоникалық толқындардың таралу процессін зерттеу проблемаларын анықтау, меншікті нысандар мен берілген пластинкалардың жиілік шектелген тербелісіне тікелей қатысты болады.

Екі қарама-қарсы шеттері топсалап бекітілген, ал басқа екі жағы әр түрлі бекіту түрлерімен ұстатылған немесе кернеу әсер етпейтіндей болып орналасқан болса, онда бұл жағдай аса күрделі тікбұрышты жазық элемент тербелісі болып табылады.

Мұндай есептер классы сандық түрде де немесе аналитикалық түрде де шешілетін меншікті жиілік тербелісін анықтайтын трансценденттік теңдеулерге акеледі

Трансценденттік жиіліктік теңдеуін алгебралық түрге келтіре отырып, тік бұрышты пластинкалар шектік шарттарын, геометриялық және механикалық сипаттағы тік бұрышты жазық элементтің тербеліс теңдеуінің әсері негізінде зерттеуге болады.

**Түйін сөздер:** пластинка, анизотропия, меншікті тербелістер, еріксіз тербелістер, трансценденттік, теңдеу, реология.

**А. Ж. Сейтмуратов<sup>1</sup>, Д. М. Жарылғапова<sup>1</sup>, Н. К. Медеубаев<sup>2</sup>, А. А. Ибраева<sup>1</sup>**

<sup>1</sup>Кызылординский государственный университет им. Коркыт Ата, Казахстан,

<sup>2</sup>Карагандинский государственный университет им. Е. Букетова, Казахстан

### **ПРИКЛАДНЫЕ ЗАДАЧИ КОЛЕБАНИЯ ПЛАСТИН ПРИ БОЛЕЕ СЛОЖНЫХ ГРАНИЧНЫХ УСЛОВИЯХ**

**Аннотация.** В настоящей работе приводятся результаты по исследованию собственных и вынужденных колебаний плоских элементов с учётом слоистости материала пластинки, реологических вязких свойств, анизотропии и т.д.

При исследовании гармонических волн в деформируемых телах вводится понятие фазовой скорости как скорости изменения состояния среды, при этом фазовая скорость выражается через частоты собственных колебаний и поэтому исследование распространения гармонических волн имеет прямое отношение к проблемам определения собственных форм и частот колебаний ограниченных в плане пластин.

Более сложным колебанием прямоугольного плоского элемента является колебание, когда два из противоположных краёв шарнирно оперты, а два других края – имеют различные виды закрепления или свободны от напряжений.

Данный класс задач приводит к трансцендентным уравнениям для определения частот собственных колебаний, которые можно решать как численно, так и аналитически.

Трансцендентные частотные уравнения можно сводить к алгебраическим и исследовать влияние, как граничных условий по краям прямоугольной пластинки или прямоугольного плоского элемента, так и параметров геометрического и механического характера на частоты собственных колебаний прямоугольных плоских элементов.

**Ключевые слова:** пластинки, анизотропия, собственные колебания, вынужденные колебания, трансцендентные, уравнения, реология.

#### **Сведение об авторах:**

Сейтмуратов Ангысын Жасаралович – д. ф.-м. н., профессор кафедры «Физика и математика» КГУ им. Коркыт Ата

Жарылғапова Дина Муратовна – к. п. н., старший преподаватель кафедры «Физика и математика» КГУ им. Коркыт Ата

Медеубаев Нурболат Куттымуратович – старший преподаватель кафедры «Алгебра, мат. логика и геометрия» КарГУ им. Букетова

Ибраева Анар Абдикаримовна – старший преподаватель кафедры «Физика и математика» КГУ им. Коркыт Ата

### **Publication Ethics and Publication Malpractice in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct ([http://publicationethics.org/files/u2/New\\_Code.pdf](http://publicationethics.org/files/u2/New_Code.pdf)). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/e4ditors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

[www.nauka-nanrk.kz](http://www.nauka-nanrk.kz)

**ISSN 2518-170X (Online), ISSN 2224-5278 (Print)**

<http://geolog-technical.kz/index.php/kz/>

Верстка *Д. Н. Калкабековой*

Подписано в печать 06.06.2017.

Формат 70x881/8. Бумага офсетная. Печать – ризограф.

18,9 п.л. Тираж 300. Заказ 3.