

ISSN 2518-170X (Online),
ISSN 2224-5278 (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
Қ. И. Сәтпаев атындағы Қазақ ұлттық техникалық зерттеу университеті

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Казакский национальный исследовательский
технический университет им. К. И. Сатпаева

NEWS

OF THE ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Kazakh national research technical university
named after K. I. Satpayev

**SERIES
OF GEOLOGY AND TECHNICAL SCIENCES**

2 (434)

MARCH – APRIL 2019

THE JOURNAL WAS FOUNDED IN 1940

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series of geology and technical sciences scientific journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of geology and technical sciences in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of geology and engineering sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Геология және техникалық ғылымдар сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді геология және техникалық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия геологии и технических наук» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК. Серия геологии и технических наук в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по геологии и техническим наукам для нашего сообщества.

Б а с р е д а к т о р ы
э. ғ. д., профессор, ҚР ҰҒА академигі

И.К. Бейсембетов

Бас редакторының орынбасары

Жолтаев Г.Ж. проф., геол.-мин. ғ. докторы

Р е д а к ц и я а л қ а с ы:

Абаканов Т.Д. проф. (Қазақстан)
Абишева З.С. проф., академик (Қазақстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Әзірбайжан)
Бакиров А.Б. проф., (Қырғыстан)
Беспәев Х.А. проф. (Қазақстан)
Бишимбаев В.К. проф., академик (Қазақстан)
Буктуков Н.С. проф., академик (Қазақстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Тәжікстан)
Грэвис Р.М. проф. (АҚШ)
Ерғалиев Г.К. проф., академик (Қазақстан)
Жуков Н.М. проф. (Қазақстан)
Қожахметов С.М. проф., академик (Қазақстан)
Конторович А.Э. проф., академик (Ресей)
Курскеев А.К. проф., академик (Қазақстан)
Курчавов А.М. проф., (Ресей)
Медеу А.Р. проф., академик (Қазақстан)
Мұхамеджанов М.А. проф., корр.-мүшесі (Қазақстан)
Нигматова С.А. проф. (Қазақстан)
Оздоев С.М. проф., академик (Қазақстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Қазақстан)
Сейтов Н.С. проф., корр.-мүшесі (Қазақстан)
Сейтмуратова Э.Ю. проф., корр.-мүшесі (Қазақстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (АҚШ)
Штейнер М. проф. (Германия)

«ҚР ҰҒА Хабарлары. Геология мен техникалық ғылымдар сериясы».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан республикасының Мәдениет пен ақпарат министрлігінің Ақпарат және мұрағат комитетінде
30.04.2010 ж. берілген №10892-Ж мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28, 219 бөл., 220, тел.: 272-13-19, 272-13-18,
<http://www.geolog-technical.kz/index.php/en/>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2019

Редакцияның Қазақстан, 050010, Алматы қ., Қабанбай батыра көш., 69а.

мекенжайы: Қ. И. Сәтбаев атындағы геология ғылымдар институты, 334 бөлме. Тел.: 291-59-38.

Типографияның мекенжайы: «Аруна» ЖК, Алматы қ., Муратбаева көш., 75.

Г л а в н ы й р е д а к т о р
д. э. н., профессор, академик НАН РК

И. К. Бейсембетов

Заместитель главного редактора

Жолтаев Г.Ж. проф., доктор геол.-мин. наук

Р е д а к ц и о н н а я к о л л е г и я:

Абаканов Т.Д. проф. (Казахстан)
Абишева З.С. проф., академик (Казахстан)
Агабеков В.Е. академик (Беларусь)
Алиев Т. проф., академик (Азербайджан)
Бакиров А.Б. проф., (Кыргызстан)
Беспаяев Х.А. проф. (Казахстан)
Бишимбаев В.К. проф., академик (Казахстан)
Буктуков Н.С. проф., академик (Казахстан)
Булат А.Ф. проф., академик (Украина)
Ганиев И.Н. проф., академик (Таджикистан)
Грэвис Р.М. проф. (США)
Ергалиев Г.К. проф., академик (Казахстан)
Жуков Н.М. проф. (Казахстан)
Кожаметов С.М. проф., академик (Казахстан)
Конторович А.Э. проф., академик (Россия)
Курскеев А.К. проф., академик (Казахстан)
Курчавов А.М. проф., (Россия)
Медеу А.Р. проф., академик (Казахстан)
Мухамеджанов М.А. проф., чл.-корр. (Казахстан)
Нигматова С.А. проф. (Казахстан)
Оздоев С.М. проф., академик (Казахстан)
Постолатий В. проф., академик (Молдова)
Ракишев Б.Р. проф., академик (Казахстан)
Сейтов Н.С. проф., чл.-корр. (Казахстан)
Сейтмуратова Э.Ю. проф., чл.-корр. (Казахстан)
Степанец В.Г. проф., (Германия)
Хамфери Дж.Д. проф. (США)
Штейнер М. проф. (Германия)

«Известия НАН РК. Серия геологии и технических наук».

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Собственник: Республиканское общественное объединение «Национальная академия наук Республики Казахстан (г. Алматы)

Свидетельство о постановке на учет периодического печатного издания в Комитете информации и архивов Министерства культуры и информации Республики Казахстан №10892-Ж, выданное 30.04.2010 г.

Периодичность: 6 раз в год

Тираж: 300 экземпляров

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28, ком. 219, 220, тел.: 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/geology-technical.kz>

© Национальная академия наук Республики Казахстан, 2019

Адрес редакции: Казахстан, 050010, г. Алматы, ул. Кабанбай батыра, 69а.

Институт геологических наук им. К. И. Сатпаева, комната 334. Тел.: 291-59-38.

Адрес типографии: ИП «Аруна», г. Алматы, ул. Муратбаева, 75

E d i t o r i n c h i e f

doctor of Economics, professor, academician of NAS RK

I. K. Beisembetov

Deputy editor in chief

Zholtayev G.Zh. prof., dr. geol-min. sc.

E d i t o r i a l b o a r d:

Abakanov T.D. prof. (Kazakhstan)
Abisheva Z.S. prof., academician (Kazakhstan)
Agabekov V.Ye. academician (Belarus)
Aliyev T. prof., academician (Azerbaijan)
Bakirov A.B. prof., (Kyrgyzstan)
Bespayev Kh.A. prof. (Kazakhstan)
Bishimbayev V.K. prof., academician (Kazakhstan)
Buktukov N.S. prof., academician (Kazakhstan)
Bulat A.F. prof., academician (Ukraine)
Ganiyev I.N. prof., academician (Tadjikistan)
Gravis R.M. prof. (USA)
Yergaliev G.K. prof., academician (Kazakhstan)
Zhukov N.M. prof. (Kazakhstan)
Kozhakhmetov S.M. prof., academician (Kazakhstan)
Kontorovich A.Ye. prof., academician (Russia)
Kurskeyev A.K. prof., academician (Kazakhstan)
Kurchavov A.M. prof., (Russia)
Medeu A.R. prof., academician (Kazakhstan)
Muhamedzhanov M.A. prof., corr. member. (Kazakhstan)
Nigmatova S.A. prof. (Kazakhstan)
Ozdoiyev S.M. prof., academician (Kazakhstan)
Postolatii V. prof., academician (Moldova)
Rakishev B.R. prof., academician (Kazakhstan)
Seitov N.S. prof., corr. member. (Kazakhstan)
Seitmuratova Ye.U. prof., corr. member. (Kazakhstan)
Stepanets V.G. prof., (Germany)
Humphery G.D. prof. (USA)
Steiner M. prof. (Germany)

News of the National Academy of Sciences of the Republic of Kazakhstan. Series of geology and technology sciences.

ISSN 2518-170X (Online),

ISSN 2224-5278 (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty)

The certificate of registration of a periodic printed publication in the Committee of information and archives of the Ministry of culture and information of the Republic of Kazakhstan N 10892-Ж, issued 30.04.2010

Periodicity: 6 times a year

Circulation: 300 copies

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19, 272-13-18,
<http://nauka-nanrk.kz/geology-technical.kz>

© National Academy of Sciences of the Republic of Kazakhstan, 2019

Editorial address: Institute of Geological Sciences named after K.I. Satpayev
69a, Kabanbai batyr str., of. 334, Almaty, 050010, Kazakhstan, tel.: 291-59-38.

Address of printing house: ST "Aruna", 75, Muratbayev str, Almaty

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN

SERIES OF GEOLOGY AND TECHNICAL SCIENCES

ISSN 2224-5278

Volume 2, Number 434 (2019), 48 – 54

<https://doi.org/10.32014/2019.2518-170X.37>

DOI 624.131.543:550.343

M. J. Ismaylov¹, S. M. Zeynalova¹, L. A. Ismaylova²

¹Institute of Geography named by acad. H. A. Aliyev of ANAS, Baku, Azerbaijan,

²Azerbaijan State Oil and Industry University, Baku, Azerbaijan.

E-mail: Miri_Ismaylov@box.az, selinsevil@gmail.com

**DYNAMICS OF THE RELIC FOREST LANDSCAPE
IN AZERBAIJAN AND WAYS TO SOLVE ENVIRONMENTAL
PROBLEMS**

Abstract. For the first time in the article, emerging the historical dynamics of the relief landscape of Azerbaijan and the ecological problems from it have been studied extensively. It has been established, plain forests completely destroyed and transposed to anthropogenic complexes; the biodiversity of existing forest ecosystems has been diminished, some species of trees and shrubs have been ruined, and the density of the forest has dropped from the ecological norm. The proposed environmental sustainability index is calculated (by the formula $\Sigma_i = S_i/S_3$) ecological stability in the studied region. Taking into consideration the role of the Relict Forest Ecosystem in the preservation of the global gene pool, the Hirkan National Park area must be expanding to three times.

Keywords: the National park Hirkan, relict forest ecosystem, initial forest, relief, natural landscape.

1. Introduction. The study of relict elements of natural complexes studies, their role in the genesis, structure, function, dynamics and evolution of landscapes is one of the important issues in modern landscape science. In the structure of Azerbaijan landscapes, it is possible to find both natural and anthropogenic relict elements. Usually, the relict elements in landscapes complicate its structure and further enhance internal diversity.

2. Objects and methods of researches. The research area is located on the west coast of the Caspian Sea in the south of the Republic of Azerbaijan. The area is 6069.0 km². The relief is located in the eastern part of the plain and in the west from the Talish, Pestasar, and Burovar mountain ranges. The absolute height of the relief ranges from 27 m to 2493 m.

Method of the research. The relief elements of the landscape are divided into two places: dynamic and static. Static relief elements include relief, lithological composition of rocks, and dynamic relict elements to fauna and flora. The research area region landscapes are mostly rich in dynamic relict elements.

The relief of the Lankaran lowland and the rocks forming it are younger than the New Caspian area. But later, the Hirkan-type relict flora and fauna species in the neighboring mountainous region migrated to the lowlands and formed existing grass-forest landscapes. Thus, the relief of the Lankaran lowland's natural forest ecosystems and the geological substrate is relatively young and the formed biocenoses have older relict elements. For this reason, modern natural landscapes of the plains are relict complexes. The forest and forest-field complexes consisting of hirkan flora formed on the volcanic sedimentary rocks of the eocene-oligocene age are the typical relict landscapes of the mountainous part of the study area [1; 9; 10].

These landscapes are the largest single relief natural system in Azerbaijan, which has reached today, leaving behind the harsh climatic conditions of millions of years of geological eras and of the fourth era of glaciation. However, during the historical periods, the natural landscapes of the Lankaran region have undergone severe structural transformation through man's economic activity and have been transformed into derivative forests in some territories (table 1).

Table 1 – Structural signs of primary and derivative (secondary) forests in Lankaran geographical area

Rich variety of plants	Age of trees	Tiers	The height of the trees	Forest structure	Growth		Age of forest	Type of plant species
					height, m	volume, m ³		
Initial forest in Astara river basin								
1000 hectare 32 species of plant	Different age composition from 30 years 100 and more than	Tiers: I-IV-tree V-bush VI-grass	Middle: from 25 to 40 m	Regular	0,26 0,40	0,14 0,45	65-80 years	Sibljakand others
Secondary forest								
Monodominant	The same age as the composition 50-60 years	2-3 yarusu I-tree II-bush III-grass	Middle: 10-15 m	Irregular	0,60 0,70	0,38 0,57	40-50 years	Thin herbs, sibljakand others

The analysis of Table 1 shows that in the natural-geographical region of Lankaran, every 1000 ha of the initial relic forest landscape were 32 types of trees and shrubs. At present, we can find relict forests, which have relatively preserved their original structure in the Hirkan National Park and the Astara river basin.

Observations made in the area near Haftoni and TazaAlvadivillages were shown the secondary forest ecosystems are more monodominant and the species of trees and bushes are 5-6 times smaller than earlier. There are no significant differences in the age of trees in such forests, just its structure is 2-3 tiers. The height of trees and the increase in wood is 1.5-2 times more than the initial forests.

Anthropogenic dynamics of forest ecosystems. As a result of man's economic activity, the area of the forest has declined, the composition has changed, the density decreases, and the natural regeneration has weakened. Currently, the relict ecosystem forest of Talish has remained in episodic form, only in the plains of the region. In the biogeographic survey there are 1167 species of plants in the Hirkan (369 km²) forest ecosystems [6]. According to this indication, Hirkan forest ecosystems occupy the first place among 18 Caucasian states of the florist. Even in the low-density subtropical Kolxida region of Georgia, the number of plant species in 1,000 hectares is three times less than the Hirkan forests.

According to [3; 11] there are 150 trees and shrubs in the dendroflora of Talish, 36 of them are endemic. Iran's Caspian region is the center of Hirkan forest ecosystems with the relic elements. The area of these forests has been more than 35,000 km².

3. Results. The major violations of the forest ecosystem with anthropogenic impacts are more common in the plain. Each landscape must fulfill a certain function in accordance with the socio-economic requirements of the existing society. The analysis shows that socio-economic functions of the forest ecosystems of Lankaran have also changed in the historical periods change, and natural complexes have been reconstructed accordingly (table 2).

Table 2 – Trends in the economic functioning of forest landscapes in Lankaran and the surrounding lowlands

Years	Area, hectares	Forest ecosystem, %	Anthropogenic landscapes, %	Trends in the economic functioning of forest landscapes
1890-1898	135 890	49 037/36	86 853/64	Rice paddy, cattle breeding, hunting
1914-1930	138 494	28 038/20	110 456/80	Rice paddy, cattle breeding, tobacco-cultivation
1930-1950	138 010	21 100/15	116 910/85	Tea-growing, rice paddy, citrus fruits growing, vegetable growing, cattle breeding
1980-2017	250 974	2618/1	248 356/99	Tea-growing, vegetable growing, cattle breeding, tourism

By the end of the 19th century, the forests were broken down for the use of rice paddy, expansion of habitat and the use of fuel. Until that time, 36% of the plain was covered with forests [2; 5]. The massive destruction of the forests was observed in the first half of the 20th century. During this period, to develop intensive tea-growing and citrus fruits, 27 937 ha of valuable relict species were destroyed by extinction. At present, up to one percent of the plains can be found with anthropogenic degradation in the forests (figure 1).




Figure 1 – Dynamics of the ratio of agro-ecosystems to ecosystems over the past 120 years in the Lankaran and surrounding lowland

Only plains of forests have been found in the territory of the Hirkan National Park. In the mountainous part of the area, there is an ancient history of deforestation [3; 6]. According to our research, the ecological potential for the initial settlement in Azerbaijan was the arid forests and forests of low mountainous and foothill, which are more accessible for human activity. These landscapes are widely spread in the middle mountainous area of Talish. The arid forests, including half-humid and humid mountain forests, have been transformed to the present-day mountain xerophyte stony and forest bushes with anthropogenic impact (figures 2, 3). Due to the destruction of forests, some species of animal perished or sharply reduced. From the analysis of historical sources has been discovered in the beginning of the XVIII century the lion (*Potheraleo*), the wild bull (*B.mstana-Zadei*) (beginning of the XIX century), Leopard (*Panthera pardus Y*) until the end of the XIX century, 1912 the white deer (*Carvuselapus Y.*), and until 1890-1932, the tiger (*Pantheratigir Y.*) was widely spread in the natural geographical region of Lankaran (table 3).

Table 3 – Dynamics of biodiversity in natural-geographical region Lankaran (M.Ismayilov, 1990)

The name of the species	Perished period, century/year
1. The tiger (<i>Panthera tigirs Y</i>)	Beginning of the XVIII century
2. The wild bull (<i>B.mstana-Zadei</i>)	Beginning of the XIX century
3. The white deer (<i>Carvuselapus Y.</i>)	1912
4. The lion (<i>Pontheraleo</i>)	1890-1932
5. The leopard (<i>Panthera pardus Y</i>)	1950

Table 4 – Distribution of available natural landscapes in Lankaran physical-geographical region

The number in map	The landscape name	Area, km ²
1	The xerophytic bush in the middle hill	938,7527
2	The forests of the low and middle mountain	1442,071724
3	Dry xerophyte steppe low mountain landscapes	491,915377
4	The meadow grass bush after the forest	712,070904
5	The low mountain steps	176,307631
6	The xerophytic steppe meadow landscapes	310,521866
7	Dry steppe meadow landscapes	305,880385
8	The hydromorphic landscapes	477,642945
9	The semi-desert of the lowland	353,127003
10	The coastal sand dune semi-desert	23,236721




Figure 2 – Natural landscapes that can be formed

The climate is one of the later changing components of the landscape from anthropogenic impacts, that is why the initial landscape map (figure 4) is based on the heat and humidity quantities (figure 4).

By the formula of Shashko (1961):

$$Md = \frac{P}{\sum(E-e)}, \tag{1}$$

P – quantities ratio of atmospheric precipitation; $\sum(E-e)$ – saturation deficit; Md – moisture supply of plants.

The advantage of this formula is expressed as the quantities ratio of atmospheric precipitation to the moisture deficiency, which is the main factor in evaporation. The obtained quantities are closely linked to the composition and productivity of plants. This index is a more reliable tool for studying humidification in complex relief areas [1; 4; 7].

The results obtained from the comparison of the above-mentioned landscape maps are shown in table 5. As is evident, the area of the Suppose forest landscapes is 4427.8 km². During the past decades, forests have been demolished and transformed into different landscapes. As a result, arid mountain forests (0.2), arid (0.0), half-humid (0.0), moisture (0.0) and Moisture-rich (0.01) forest landscapes. Accordingly, xerophyte landscapes were largely transformed into dry steppe and grassy landscapes.




Figure 3 – Natural landscapes available at present

Role of forest ecosystems in the protection of ecological stability. For the sustainable development of the region is important to explore the environmental sustainability and the ways to optimize.

For this purpose, the role of forest ecosystems has been assessed in environmental sustainability. Currently, the total area of forests is 1472.3 km² in the Lankaran natural-geographical area, which makes up of the total area 24.3%. At present, the ecological situation of the region is closely linked to the structural and functional characteristics of forest ecosystems. We have identified forest covering and environmental sustainability indexes in the study area.

The forest cover index is calculated by the following formula[1; 7; 12]:

$$Mi = S_1/S_2 ,$$

Mi – forest coverage index; S₁ – forest covered area; S₂ – a general comparison area.

The environmental sustainability index was calculated by the ratio of forest covered area to the total area of agro-ecosystems:

$$\Sigma i = S_1/S_3 ,$$

here Σi – the index of ecological stability; S₃ – total area of comparable agroecosystems.

Table 5 – Comparative analysis of suppose landscapes and existing natural landscape in natural humidification conditions

#	Moisture supply of plants, $Md = \frac{P}{\Sigma(E-e)}$	Moisture	Suppose landscapes	Area, km ² /% (S ₁)	Existing natural landscape	Area of the suppose landscapes, km ² /% (S ₂)	The transformation indexes of landscapes $T_i = S_2/S_1$
1	<0,10	Arid	Semi-desert	$\frac{288,4}{5,1}$	Hydromorphic	277,3	0,3
					Semi-desert	74,1	
2	0,10-0,15	Semi-arid	Dry plain	$\frac{670,7}{12,1}$	Semi-desert	279,0	0,4
					Dry steppe	291,7	
3	0,15-0,25	Arid	Dry xerophyte steppe low mountain	$\frac{219,5}{3,9}$	Dry steppe	183,4	0,5
					Xerophyte steppe	108,4	
4	0,25-0,35	Semi-arid	a. Arid forests in low and middle mountains	$\frac{159,3}{28,5}$	Xerophyte bush	938,8	0,2
					Dry steppe	347,2	
					Arid forest	312,3	
			4 b. Arid forests landscapes of plains	$\frac{387,6}{6,9}$	Dry steppe	239,9	0,0
Hydromorphic	147,7						
5	0,35-0,45	Semi-humid	5a. Semi-humid mountain forests landscapes	$\frac{109,7}{19,6}$	Dry xerophyte steppe	498,6	0,5
					Mountain-forest	600,1	
			5b. Semi-humid forest landscapes of plains	$\frac{152,4}{2,7}$	Meadow-bush	152,4	0,0
6	0,45-0,60	Humid	6a. Humid forests in low mountain	$\frac{264,5}{2,6}$	Mountain-forest	264,5	1,0
					6b. Humid forests landscapes of plains	$\frac{185,6}{3,3}$	Semi-desert
			Dry bush	162,4			
7	>0,60	Moisture-rich	Moisture-rich forests of plains and mountains	$\frac{740,7}{13,2}$	Meadow-bush	732,9	0,01
					Moisture-rich forests	7,8	

4. Conclusion.

1. The horizontal and vertical structure of the Talish forest ecosystems has been seriously violated in the past 100-150 years. The forests have been almost destroyed and transformed into anthropogenic complexes. The existing mountain forest ecosystems have diminished biodiversity, some species of trees have been severely damaged, and the density of the forest has dropped from the ecological norm.

2. For the first time, the environmental sustainability index was calculated by the formula $\Sigma i = S_1/S_3$, and it was determined that the environmental sustainability index for the region was above the average country level and equaled to 0.5 in the study area.

3. Taking into account the role of the region in the preservation of the unique genetic background of the unique relict forest landscape, the Hirkan National Park is to be expanded three times, creating relatively small areas of special protection requiring flexible management.

М. Дж. Исмаилов¹, Зейналова С. М.¹, Л. А. Исмаилова²

¹ӘҒҒА География институты, Баку, Әзірбайжан,

²Әзірбайжан мемлекеттік Мұнай және өнеркәсіп университеті, Баку, Әзірбайжан

**ӘЗІРБАЙЖАННЫҢ ҚАЛДЫҚ ОРМАН ЛАНДШАФТЫНЫҢ ДИНАМИКАСЫ
ЖӘНЕ ОРМАНДАРДЫҢ ЭКОЛОГИЯЛЫҚ МӘСЕЛЕЛЕРІН
ШЕШУ ЖОЛДАРЫ**

М. Дж. Исмаилов¹, Зейналова С. М.¹, Л. А. Исмаилова²

¹Институт Географии НАНА, Баку, Азербайджан,

²Азербайджанский государственный университет Нефти и промышленности, Баку, Азербайджан

ДИНАМИКА РЕЛИКТОГО ЛЕСНОГО ЛАНДШАФТА АЗЕРБАЙДЖАНА И ПУТИ РЕШЕНИЯ ЭКОЛОГИЧЕСКИХ ПРОБЛЕМ ЛЕСОВ

Аннотация. Впервые в статье была изучена историческая динамика реликтового ландшафта Азербайджана. Реликтовые ландшафты – сохранившиеся до настоящего времени черты различных компонентов древних ландшафтов, которые вырабатывались при иных физико-геогр. условиях и представляют собой наследие былых природных условий. Б. Б. Польшов (1925) отметил, что в каждом ландшафте и почве наряду с консервативными (современными) и реликтовыми элементами следует различать и прогрессивные элементы, характеризующие тренд дальнейшего развития. При изучении реликтовых ландшафтов необходимо учитывать реликтовый характер всех компонентов, слагающих реликтовый ландшафтный комплекс и отдельные его компоненты. В первом случае набор реликтовых ландшафтов достаточно узок, а во втором – необычайно широк.

В статье было установлено, что нами изученные равнинные леса (Лянкяранская равнина) полностью разрушены и трансформированы в антропогенные комплексы, биоразнообразие существующих лесных экосистем уменьшено, некоторые виды деревьев и кустарников разрушены, а плотность леса снизилась с экологической нормы. В первые нами предложенной формулой $\Sigma_i = S_1/S_3$ проведен расчет индекса экологической устойчивости ландшафтов в регионе.

Ключевые слова: реликтовые ландшафты, трансформация, фактор, экологический индекс.

Information about authors:

Ismaylov Mirnukh Javadoglu, candidate of geographical sciences, associate professor, leading research worker of Institute of Geography named by acad. H. A. Aliyev of ANAS, Baku, Azerbaijan; Miri_Ismaylov@box.az

Zeynalova Saida Madatgyzi, candidate of geographical sciences, associate professor, leading research worker of Institute of Geography named by acad. H. A. Aliyev of ANAS, Baku, Azerbaijan; selinsevil@gmail.com

Ismaylova Latifa Arifgyizi, Ph.D. student, Azerbaijan State Oil and Industry University, Baku, Azerbaijan

REFERENCES

- [1] Lee S. (2005). Application of logistic recreation model and its validation for landslide susceptibility mapping using GIS and remote sensing data // *International Journal of remote sensing*. 26 (7). P. 1477-1991.
- [2] Bintanja R., van de Wal R.S.W. (2008). North American ice-sheet dynamics and the onset of the 100,000-year glacial cycles // *Nature*. 454. P. 869-872.
- [3] Beruchashvili N.L. (1979). Landscape Map of Caucasus. T, scale 1:1,000,000. Ecoregion Conservation Plan for the Caucasus // 2012 revised and updated edition. WWF.
- [4] Thompson J.A., Bell J.C., Butler C.A. (2001). Digital elevation model resolution: effects on terrain attribute calculation and quantitative soil-landscape modeling // *Geoderma*. 100. P. 67-89.
- [5] Zhilin Li, Qing Zhu, Chris Gold. (2005). Digital terrain modeling: principles and methodology. CRC PRESS. Boca Raton London New York Washington, D.C. 318 p.
- [6] Garibov Y., Mardanov I., Ismaylova N., Ahmadova G., Aliyeva R. (2016). Ekzogenetic features landscapes of high mountains of the Greater Caucasus with in Azerbaijan Republic // *British Journal of Educational and Scientific Studies, Source Normalized Impact per Paper (SNIP): 5.796, SCImago Journal Rank (SJR): 5.925. N 1(23). January - June, 2016. Vol. XII. "Imperial College Press". P. 579-585.*
- [7] Kristensen T.B., Huuse M., Piotrowski J.A., Clausen O.R. (2007). A morphometric analysis of tunnel valleys in the eastern North Sea based on 3D seismic data // *Journal of Quaternary Science*. 22(8). 801-Korte M., Constable C.G. 2005.
- [8] Gerasimov I.P. (1981). Structure géologique et relief du Caucase // *Revue de géographie alpine*. 69/2. P. 225-39.
- [9] Jacques D. (1995). The Rise of Cultural Landscapes // *International Journal of Heritage Studies*. 1-2. 91-101.
- [10] Rössler M. (2006). World Heritage Cultural Landscapes // *Landscape Research*. 31: 4. P. 333-353.
- [11] James Solomon, Tatyana Shulkina, George E Schatz and st. (2014). Red List of the Endemic Plants of the Caucasus // Missouri Botanical Garden Press. Vol. 125.
- [12] Peter Goldblatt, Dale E Johnson. (2006). Index to Plant Chromosome Numbers. 2001-2003. Missouri Botanical Garden Press. Vol. 106.

**Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan**

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

Правила оформления статьи для публикации в журнале смотреть на сайте:

www.nauka-nanrk.kz

ISSN 2518-170X (Online), ISSN 2224-5278 (Print)

<http://www.geolog-technical.kz/index.php/en/>

Верстка Д. Н. Калкабековой

Подписано в печать 12.04.2019.

Формат 70x881/8. Бумага офсетная. Печать – ризограф.
15,2 п.л. Тираж 300. Заказ 2.